

Topology-Free Type Structures with Conditioning Events*

Pierfrancesco Guarino[†]

1.IX.2023

Revise & Resubmit at “Economic Theory”

Abstract

Starting without any topological assumption, we establish the existence of the universal type structure in presence of—possibly uncountably many and topologically unrestricted—conditioning events, namely, a type structure that is non-redundant, belief-complete, terminal, and unique up to measurable type isomorphism, by performing a construction in the spirit of the hierarchical one in Heifetz & Samet (1998). In particular, we obtain the result by exploiting arguments from category theory and the theory of coalgebras, thus, making explicit the mathematical structure underlying all the constructions of large interactive structures and obtaining the belief-completeness of the structure (unattainable via the standard hierarchical construction *à la* Heifetz & Samet (1998)) as an immediate corollary of known results from these fields. Additionally, we show how our construction, with its lack of topological and cardinality assumptions on the family of conditioning events, can be employed in various game-theoretical contexts.

Keywords: Universal Type Structure, Non-Redundancy, Belief-Completeness, Terminality & Uniqueness, Conditional Probability Systems, Topology-Free, Category Theory & Coalgebras.
JEL Classification Number: D80, D82.

“Before category theory was invented it had been noted in some parts of mathematics that from some privileged objects, often called ‘universal’, one could construct all objects of the same kind.”

—Dieudonné (1989, Part 1.IV.8.C, p.151)

*The present work is a considerable development of Guarino (2017), where a sketch of a proof of the result obtained with radically different tools was presented. The main technical bulk of this work has been obtained during an informal visiting period at LUISS Guido Carli back in the early spring of 2018 for which I am extremely grateful to Marco Scarsini. I would like to thank the editor, Nicholas Yannellis, the anonymous handling editor, and two referees for their outstanding reading of the paper and their suggestions that ultimately lead to the present version. Special thanks go to Pierpaolo Battigalli, for his long-term interest in this endeavour, and to Martin Meier, without whose encouragement to pursue the project—quite simply—this manuscript would have never become public. I also would like to thank Michael Greinecker and Gabriel Ziegler for many—most stimulating—conversations on the topic of large interactive structures in general and Marcus Pivato for enlightening discussions on category theory. Finally, I would like to thank the audiences of the EWET 2023 Workshop and the SAET 2023 Conference. Of course, all errors are mine. I thankfully acknowledge financial support from the Austrian Science Fund (FWF) (P31248-G27) and from MIUR under the PRIN 2017 program (grant number 2017K8ANN4).

[†]University of Udine (Department of Economics and Statistics – DIES). *E-mails:* pf.guarino@hotmail.com & pierfrancesco.guarino@uniud.it.

CONTENTS

1	Introduction	3
1.1	Motivation & Results	3
1.2	Related Literature	5
1.3	Synopsis	5
2	Mathematical Preliminaries	6
2.1	Measure Theory	6
2.2	Theory of Coalgebras	8
3	Type Structures	10
3.1	Basic Definition	10
3.2	Relation Between Type Structures	10
3.3	Infinite Hierarchies of Beliefs	11
3.4	Large Type Structures	12
4	Universality	13
4.1	There (in the Coalgebraic Framework)...	13
4.1.1	Obtaining the Terminal Carriers	15
4.1.2	Obtaining the Terminal Transition	16
4.1.3	Establishing Terminality	19
4.2	...In the Dale (between the Two Worlds)...	20
4.3	...and Back Again (to Type Structures)	22
5	Applications	23
5.1	Topology-Free Conditioning Events	24
5.2	Uncountability of the Conditioning Events	24
5.3	Nature of the Conditioning Events	25
6	Discussion	26
6.1	Relation to the Literature	26
6.1.1	Relation to Heifetz & Samet (1998)	26
6.1.2	Relation to Viglizzo (2005b)	26
6.1.3	Relation to Heinsalu (2014)	27
6.2	Relation to Universality Notions	27
6.2.1	On the Notion of Kolmogorov's Consistency	27
6.2.2	On Belief-Completeness & Relation to Meier (2012)	28
6.2.3	On the Notion of Universality	29
6.3	On Technical Aspects of the Construction	30
6.3.1	On Type Structures	30
6.3.2	On Type Morphisms	30
6.3.3	On the Family of Conditioning Events	30
A	Category Theory Background	31
B	Measure-Theoretic Results	33
C	Proofs of Section 4	34
	References	39

1. INTRODUCTION

1.1 Motivation & Results

Type structures are one of the most widely employed tools in theoretical economics. Introduced in [Harsanyi \(1967\)](#) to handle the technical problems arising from the infinite regress proper of the analysis of games with incomplete information, their usage ranges from applied issues, such as those tackled by mechanism design, to more foundational problems, as exemplified by those works belonging to epistemic game theory.¹

In the context of type structures, a particular role is played by those type structures that satisfy what can be deemed ‘large’ properties, which are typically captured via the so-called notion of universality.² Intuitively, the Universal Type Structure, where the usage of the definite article comes from the fact that the construct is essentially *unique*, is a particular type structure that contains all the possible types that can arise given a certain domain of uncertainty. Thus, it is immediate to see the importance of this notion in the fields mentioned above, as—for example—pointed out in [Bergemann & Morris \(2012, Section 3\)](#): employing the Universal Type Structure allows the analyst to drop all at once the possible *a priori* assumptions that can be forced upon a game-theoretical analysis by employing smaller type structures, since, in the evocative words of [Aumann & Heifetz \(2002, Section 8, pp.1672–1673\)](#),

“The system does not depend on reality; it is a framework, it fits any reality, so to speak, like the frames that one buys in photo shops, which do not depend on who is in the photo and they fit any photo with any subject, as long as the size is right. In brief, there is no substantive information in the system.”

In other words, the usage of smaller type structures forces upon a game-theoretical analysis certain restrictions by *a fortiori* ruling out some possible beliefs, i.e., contrary to the universal one, there is “*substantive information in the system*” when such a system is a ‘small’ type structure—even if the substantive information is present somewhat implicitly.

Thus, working with the Universal Type Structure is a considerable step further in dropping involuntary—and non-explicit—assumptions that could limit game-theoretical analysis: as such, employing the Universal Type Structure is—for example—a typical (even if admittedly extreme) step when an analyst wants to relax common knowledge assumptions in the spirit of the so-called *Wilson’s doctrine* as in [Wilson \(1987\)](#). However, there is another way in which potential non-explicit assumptions could arise even when using the Universal Type Structure, namely, when the Universal Type Structure is built from topological assumptions. Indeed, in such a case, it is most natural to ask ourselves if the results we obtain from a given analysis are the consequence of the very topological assumptions needed to build the Universal Type Structure employed. Thus, it is in this context that topology-free constructions of Universal Type Structures happen to derive their importance: they are constructs that can be used to perform analyses robust to topological details and can provide a conceptual foundation for the topological constructions.

For the ‘standard’ case appropriate to deal with—for example—static games, the construction of the Universal Type Structure has been obtained in [Heifetz & Samet \(1998\)](#). Rather crucially, this work did not follow the standard procedure based on the construction of coherent infinite hierarchies of beliefs (i.e., infinite hierarchies of beliefs satisfying a coherency requirement across different belief orders) as performed when starting from topological assumptions, in the spirit of [Mertens & Zamir \(1985\)](#) or [Brandenburger & Dekel \(1993\)](#).³ Indeed, this was a consequence of the striking result obtained in [Heifetz & Samet \(1999\)](#), who showed that, when we drop topological assumptions, we lose the equivalence between types (as coherent infinite hierarchies of beliefs that admit an extension to the limit) and coherent infinite hierarchies of beliefs and there are coherent infinite hierarchies of beliefs that are actually *not* types. As a consequence, the approach taken in [Heifetz & Samet \(1998\)](#), based on taking types belonging to ‘small’ type structures as ready-made objects to then collect all of them in a large type structure, had an undesirable side-effect. That is, the obtained construct had two out of the three properties attached to the idea of universality,

¹Regarding epistemic game theory, see the survey [Dekel & Siniscalchi \(2015\)](#), or the two textbooks [Perea \(2012\)](#) and [Battigalli et al. \(Work in Progress\)](#), completely devoted to the topic.

²See [Section 3.4](#) for the definition we employ in this work and [Section 6.2.3](#) with respect to how the term is used in the literature.

³See also [Armbruster & Böge \(1979\)](#), [Heifetz \(1993\)](#), and [Mertens et al. \(2015, Chapter A.III.1\)](#).

namely, terminality (capturing the very idea of collecting all those types in the same large structure) and non-redundancy: the remaining property, namely, belief-completeness, particularly desirable for epistemic endeavours, had to be established in Meier (2012) by employing tools from infinitary probability logic.⁴

In the present work, we build the Universal Type Structure with conditioning events without topological assumptions, thus, extending the work of Heifetz & Samet (1998) to the presence of conditioning events and the work of Battigalli & Siniscalchi (1999), that constructed the Universal Type Structure *à la* Brandenburger & Dekel (1993) starting from a Polish common domain of uncertainty to deal with the presence of conditioning events, to a topology-free setting. Incidentally, by obtaining our result, we prove true a conjecture made in Battigalli & Siniscalchi (1999, Section 2, p.198) regarding the possibility of performing a construction *à la* Heifetz & Samet (1998) for type structures with conditioning events. However, whereas to obtain the result we follow the construction performed in Heifetz & Samet (1998), at the same time, we actually take a more general approach, relying on tools from category theory and the theory of coalgebras (duly introduced in Section 2.2). In taking this path, built on work by Viglizzo (2005b),⁵ where a coalgebraic analog of Heifetz & Samet (1998) for the case without conditioning events is derived, we obtain in Theorem 1 all at once the three universality properties of our resulting construct, which is—indeed—non-redundant, belief-complete, and terminal (and unique up to—measurable—type isomorphism).⁶

With respect to the tools chosen to obtain our result, it is important to stress one point: there is no ‘shortcut’ to get *all* the relevant universality properties at once in the topology-free case via a construction that does not involve a certain level of ‘technicalities’. Indeed, as pointed out above, it is the construction in Meier (2012) that actually established for the first time the universality of the construction in Heifetz & Samet (1998) by—among other things—proving its belief-completeness by employing tools from infinitary probability logic, with Viglizzo (2005b) being an alternative based on category theoretical and coalgebraic tools. Thus, in order to establish the universality in the topology-free case, either we choose Scylla or Charybdis, i.e., either we pick tools from infinitary probability logic or we choose the category theoretical and coalgebraic approach:⁷ the standard construction of Heifetz & Samet (1998) does not *automatically* deliver everything we would like to get with belief-completeness being the missing ingredient.⁸

Thus, to establish this result, we employ the following strategy. First of all, in Section 4.1, we bypass the presence of interactive agents and we show that there is essentially a basic mathematical structure underlying the very notion of type structure (in presence of conditioning events), namely, a coalgebra (as in Definition 2.5). As a result, we establish the existence of a terminal coalgebra in Proposition 2. Armed with this result, in Section 4.2, we tackle the presence of interactive agents and we show that the main bulk of the endeavour has already been established, since a minimal change allows us to obtain Proposition 3, which is a translation of Theorem 1 in coalgebraic terms as set in Proposition 2, with the caveat that we now take care of the presence of interactive agents. Finally, in Section 4.3, we show that type structures are indeed coalgebras as defined in Section 4.2, thus, exploiting Proposition 3 to establish Theorem 1 essentially as its immediate corollary. It is important to underline that, to proceed along the lines just described, we closely follow the path and the proofs in Viglizzo (2005b) (reproduced here with our notation for self-containment purposes), in themselves—as already mentioned—a translation in coalgebraic terms of those in Heifetz & Samet (1998). Hence, from a technical standpoint, the crucial innovation of this paper lies in identifying the specific properties of the mathematical constructs we have to work with (i.e., product conditional measurable spaces sharing a family of conditioning events as in Definition 2.3), a point which has an immediate impact on the peculiarities of the categorical construct (i.e., the functor) we have to work with and on the proofs of all these results that explicitly refer to the presence of conditioning events, that are the ‘new’ ones with respect to Viglizzo (2005b).

⁴See Section 6.2.2 for a discussion of the result and the path chosen to establish it.

⁵See Lawvere (1962) and Giry (1982) for the first works addressing measure and probability spaces from a categorical standpoint.

⁶See Section 6.2.3 regarding the notion of universality we employ and its relation to terminality and the other notions mentioned above.

⁷With the understanding that the disjunction is used here in an inclusive sense: i.e., it is possible to work with *coalgebraic modal logic*. See Section 6.2.2 for a discussion of this point.

⁸Where, in particular, what is missing is that the belief functions are *isomorphisms* in the appropriate category under scrutiny. See Section 6.2.3 concerning this point.

It is important to stress that we see this work as having *two* main applications. The first concerns the result itself and how it can be used in game theory. Indeed, whereas in [Section 5](#) we delve into these issues by providing examples of where our result can prove to be useful for game-theoretical endeavours, here we briefly point out some immediate contexts where it can be employed. Thus, for example, beyond being the reference framework for [Bergemann & Morris \(2005, Section 2.5\)](#), it should be observed that the construct obtained in [Heifetz & Samet \(1998\)](#) has been used to obtain the epistemic characterization of Interim Correlated Rationalizability as in [Dekel et al. \(2007, Section 4.1\)](#) and to study Rationalizability in games with incomplete information in general as in [Battigalli et al. \(2011, Section 1.3\)](#). Thus, [Theorem 1](#) (and the construction obtained here more in general) can provide the ground for similar works—in the spirit of [Penta \(2015\)](#) and [Müller \(2016\)](#)—where there is the need to drop topological assumptions by also taking into account the presence of conditioning events possibly without any restriction, either topological or cardinal, on the nature of them. The second application, that we consider as important and potentially far-reaching as the previous one, concerns the introduction of the tools we employ to an ‘economics’ audience. Indeed, whereas category theory and the theory of coalgebras can be used *directly* to tackle specific technical and conceptual problems belonging to economic theory, as we do here and it has already been done in works referenced in [Section 1.2](#), those fields can both prove to be extremely useful in a more indirect way. To see this, first of all, it is important to recognize, in the words of [Leinster \(2014, Introduction, p.1\)](#), that

“Category theory takes a bird’s eye view of mathematics. From high in the sky, details become invisible, but we can spot patterns that were impossible to detect from ground level.”

Thus, it is possible to use category theory in an indirect—heuristic—fashion. Here, we refer to the fact that being acquainted to the categorical language and its results could in principle allow to see patterns “*from high in the sky*”; this, in turn, could allow to conjecture certain solutions to specific problems proper of economic theory exactly via the fact that those solutions would be proper of more general settings, as showed via results in category theory. In other words, observing that something is true for many different categories all sharing certain ‘nice’ properties could prove to be useful to deal with specific problems, once the right category to address these problems has been identified and it turns out that this category has those very same ‘nice’ properties. And here one interesting point is in order: this heuristic process should not ask to write results using the categorical language, which could potentially remain in the background.

1.2 Related Literature

This work is related to various streams of literature. Of course, it is related to the literature focusing on the existence of large type structure, most notably [Heifetz & Samet \(1998\)](#), in working in a topology-free setting, [Battigalli & Siniscalchi \(1999\)](#), in addressing the presence of conditioning events, [Meier \(2012\)](#), in dealing with the problem of belief-completeness, and [Viglizzo \(2005b\)](#), for the tools employed. In using machinery from category theory, it is related to an emerging literature in economic theory that employs these tools, such as [Heinsalu \(2014\)](#), [de Oliveira \(2018\)](#), [Galeazzi & Marti \(2023\)](#), and [Pivato \(Work in Progress\)](#). Also, regarding the usage of coalgebras,⁹ it is related to [Moss & Viglizzo \(2004\)](#) and—in particular, once more—[Viglizzo \(2005b\)](#),¹⁰ along with the aforementioned [Heinsalu \(2014\)](#) and [Galeazzi & Marti \(2023\)](#), and unpublished work by Davide Ferri.¹¹

1.3 Synopsis

This paper is organized as follows. In [Section 2](#), we introduce the mathematical notions from measure theory and the theory of coalgebras we need for our endeavour. In [Section 3](#), we introduce type structures along with related notions. In [Section 4](#), we prove the existence of the topology-free universal type structure with conditioning events, while in [Section 5](#) we show various contexts where

⁹[Heifetz \(1996\)](#) is a paper that should be—indirectly—related to this stream of literature via its relation with [Aczel \(1988\)](#), where it is possible to find in Chapter 7 an explicit construction such as the one in [Section 4.1](#).

¹⁰See also [Viglizzo \(2005a\)](#), [Moss & Viglizzo \(2006\)](#), and [Moss \(2011\)](#).

¹¹As an M.Sc. thesis at Bocconi University, starting from topological assumptions (in particular, starting from a Polish common domain of uncertainty as in [Battigalli & Siniscalchi \(1999\)](#)—personal communication).

this object can be used. Finally, in [Section 6](#), we address some points related to our result and we discuss the relation between the present work and various works belonging to the same stream of literature. Regarding the appendices, we devote [Appendix A](#) to a self-contained introduction to the notions from category theory needed for our purposes, we collect results from measure theory we employ to establish the universality in [Appendix B](#), whereas we relegate to [Appendix C](#) all the proofs of our results that cannot be found in the main body of this work.

2. MATHEMATICAL PRELIMINARIES

2.1 Measure Theory

Let (M, Σ_M) be a measurable space, that is, a set M endowed with a σ -algebra Σ_M : in the following, whenever we refer to a set M as a measurable space without explicitly denoting its σ -algebra, it is assumed that M is endowed with a σ -algebra Σ_M . Given a measurable space (M, Σ_M) , the σ -algebra Σ_M is *separative* if for every $x, x' \in M$ with $x \neq x'$ there exists an $E_x \in \Sigma_M$ such that $x \in E_x$ and $x' \notin E_x$. We let every (finite or countably infinite) product of measurable spaces be endowed with the product σ -algebra,¹² i.e., given an arbitrary product space $M := \prod_{\lambda \in \Lambda} M_\lambda$ with $(M_\lambda, \Sigma_\lambda)$ measurable and π_λ denoting the projection function as canonically defined over the index set Λ , for every $\lambda \in \Lambda$, we endow M with the *product σ -algebra*

$$\bigotimes_{\lambda \in \Lambda} \Sigma_\lambda := \sigma(\{ \pi_\lambda^{-1}(E) \mid \lambda \in \Lambda, E \in \Sigma_\lambda \}),$$

where we—alternatively—denote this σ -algebra with $\bigotimes_{\lambda \in \Lambda} \Sigma_\lambda$ (as we do above) or $\Sigma_{\prod_{\lambda \in \Lambda} M_\lambda}$. Given a measurable space $(M', \Sigma_{M'})$, a set M , and a function $f \in M'^M$, we let $\sigma(f)$ denote the smallest σ -algebra on M that makes f measurable. A *measurable isomorphism* between two measurable spaces (M, Σ_M) and $(M', \Sigma_{M'})$ is a bijective measurable function $f \in M'^M$ such that $f(E) \in \Sigma_{M'}$ for every $E \in \Sigma_M$ and $f^{-1}(E') \in \Sigma_M$ for every $E' \in \Sigma_{M'}$, where—to lighten the notation—we typically omit the reference to the σ -algebras that make a function measurable. Also, given an arbitrary set M , we let id_M denote the identity function on M , which is—trivially—a measurable isomorphism whenever M is a measurable space.

We let $\Delta(M)$ denote the set of all σ -additive probability measures over M (henceforth, probability measures). Given a measurable space (M, Σ_M) and a possibly uncountable¹³ subset $\mathcal{C}_M \subseteq \Sigma_M \setminus \{\emptyset\}$ of *conditioning events*, we call the space $(M, \Sigma_M, \mathcal{C}_M)$ a *conditional measurable space*.

Definition 2.1 (Conditional Probability System). A conditional probability system¹⁴ (henceforth, CPS) on a conditional measurable space $(M, \Sigma_M, \mathcal{C}_M)$ is a function

$$\nu(\cdot|\cdot) : \Sigma_M \times \mathcal{C}_M \rightarrow [0, 1]$$

that satisfies the following axioms:

C1. For every $C \in \mathcal{C}_M$, $\nu(C|C) = 1$;

C2. For every $C \in \mathcal{C}_M$, $\nu(\cdot|C) \in \Delta(M)$;

C3. For every $E \in \Sigma_M$ and $D, C \in \mathcal{C}_M$, if $E \subseteq D \subseteq C$, then $\nu(E|C) = \nu(E|D) \cdot \nu(D|C)$.

We let $[\Delta(M)]^{\mathcal{C}_M}$ denote the set of all functions from \mathcal{C}_M to $\Delta(M)$, while we let $\Delta^{\mathcal{C}_M}(M) \subseteq [\Delta(M)]^{\mathcal{C}_M}$ denote the set of CPSs on $(M, \Sigma_M, \mathcal{C}_M)$, with $\nu := (\nu(\cdot|C))_{C \in \mathcal{C}_M} \in \Delta^{\mathcal{C}_M}(M)$. We let

$$\gamma_C^p(E) := \{ \nu \in \Delta^{\mathcal{C}_M}(M) \mid \nu(E|C) \geq p \},$$

for every $E \in \Sigma_M$, $C \in \mathcal{C}_M$, and $p \in [0, 1]$.

¹²See [Srivastava \(1998, Chapter 3.1, p.87\)](#).

¹³See [Section 6.3.3](#) for a discussion of this point.

¹⁴This corresponds to one of the primitive elements of a *conditional probability space* of [Rényi \(1955, Sections 1.2 & 1.4\)](#). However, the name—that eventually stuck in the game-theoretic literature—actually comes from a related definition from [Myerson \(1986, Section 5, pp.336–337\)](#). See [Hammond \(1994, Section 3.2\)](#) for a discussion of the relation between these two notions.

Definition 2.2 (σ -Algebra on the Space of CPSs). Given a conditional measurable space $(M, \Sigma_M, \mathcal{C}_M)$, the space $\Delta^{\mathcal{C}_M}(M)$ is endowed with the σ -algebra¹⁵

$$\Sigma_{\Delta^{\mathcal{C}_M}(M)} := \sigma(\{ \gamma_C^p(E) \mid E \in \Sigma_M, C \in \mathcal{C}_M, p \in [0, 1] \}). \quad (2.1)$$

Given a conditional measurable space $(M, \Sigma_M, \mathcal{C}_M)$, in this paper, we focus on a specific family of spaces tightly linked to $(M, \Sigma_M, \mathcal{C}_M)$, introduced with the idea of finding a ‘common’ family of exogenously imposed conditioning events between product spaces, that we define next.¹⁶

Definition 2.3 (Product Conditional Measurable Spaces Induced by Conditioning Events). Given a conditional measurable space $(M, \Sigma_M, \mathcal{C}_M)$ and a measurable space (X, Σ_X) , the space $(M \times X, \Sigma_M \otimes \Sigma_X, \mathcal{C}_{M \times X})$ is a product conditional measurable space induced by $(M, \Sigma_M, \mathcal{C}_M)$ if

$$\mathcal{C}_{M \times X} := \{ D_X \in \Sigma_M \otimes \Sigma_X \mid \exists C \in \mathcal{C}_M : D_X := C \times X \}. \quad (2.2)$$

When it does not lead to any ambiguity, we drop from [Definition 2.3](#) the reference to the family of conditioning events and we simply use the expression “product conditional measurable space”. Also, two product conditional measurable spaces $(M \times X, \Sigma_M \otimes \Sigma_X, \mathcal{C}_{M \times X})$ and $(M \times Y, \Sigma_M \otimes \Sigma_Y, \mathcal{C}_{M \times Y})$ induced by \mathcal{C}_M are said to *share* \mathcal{C}_M . Now, observing the structure of $\mathcal{C}_{M \times X}$ in [Equation \(2.2\)](#), we have that the conditioning events of spaces sharing \mathcal{C}_M are always induced from the conditioning events in \mathcal{C}_M , which allows us to introduce the following notational conventions extensively employed throughout this work.

Notation 1 (Common Conditioning Events). Given a conditional measurable space $(M, \Sigma_M, \mathcal{C}_M)$, for every product conditional measurable space $(M \times X, \Sigma_M \otimes \Sigma_X, \mathcal{C}_{M \times X})$ induced by \mathcal{C}_M , we write:

- $(M \times X, \Sigma_M \otimes \Sigma_X, \mathcal{C}_M)$ instead of $(M \times X, \Sigma_M \otimes \Sigma_X, \mathcal{C}_{M \times X})$,
- $\Delta^{\mathcal{C}_M}(M \times X)$ instead of $\Delta^{\mathcal{C}_{M \times X}}(M \times X)$, and
- $(\nu(\cdot|C))_{C \in \mathcal{C}_M} \in \Delta^{\mathcal{C}_M}(M \times X)$ instead of $(\nu(\cdot|C \times X))_{C \in \mathcal{C}_M}$.

In particular, whenever we use the notation $\Delta^{\mathcal{C}_M}(M \times X)$, it is understood that $M \times X$ is induced by \mathcal{C}_M , i.e., $\mathcal{C}_{M \times X}$ is defined as in [Equation \(2.2\)](#).

Given a conditional measurable space $(M, \Sigma_M, \mathcal{C}_M)$, two product conditional measurable spaces $(M \times X, \Sigma_M \otimes \Sigma_X, \mathcal{C}_{M \times X})$ and $(M \times Y, \Sigma_M \otimes \Sigma_Y, \mathcal{C}_{M \times Y})$ sharing \mathcal{C}_M , and a measurable function $f \in (M \times Y)^{M \times X}$, the *image measure with conditioning events* of f is the function

$$\widehat{f} := (\widehat{f}_{D_Y})_{D_Y \in \mathcal{C}_{M \times Y}} : \Delta^{\mathcal{C}_{M \times X}}(M \times X) \rightarrow \Delta^{\mathcal{C}_{M \times Y}}(M \times Y),$$

defined as

$$\widehat{f}_{D_Y}(\nu)(E) := \nu(f^{-1}(E)|D_X),$$

for every $D_Y \in \mathcal{C}_{M \times Y}$ and $D_X \in \mathcal{C}_{M \times X}$ such that there exists a $C \in \mathcal{C}_M$ with $\pi_M D_Y = \pi_M D_X = C$, CPS $\nu := (\nu(\cdot|D_X))_{D_X \in \mathcal{C}_{M \times X}} \in \Delta^{\mathcal{C}_{M \times X}}(M \times X)$, and $E \in \Sigma_M \otimes \Sigma_Y$.¹⁷ We rephrase for future reference what we just introduced in the definition that follows, which exploits the notational conventions set forth in [Notation 1](#).

¹⁵When M is Polish and $\Delta(M)$ is endowed with the topology of weak convergence, this σ -algebra (modulo presence of conditioning events) is exactly the Borel σ -algebra induced by the topology of weak convergence (see [Kechris \(1995, Theorem 17.24, p.112\)](#)). Additionally, when in presence of \mathcal{C}_M , with this σ -algebra we have that $\Delta^{\mathcal{C}_M}(M)$ is closed in $[\Delta(M)]^{\mathcal{C}_M}$, as established in [Battigalli & Siniscalchi \(1999, Lemma 1, p.193\)](#) (see also [Battigalli & Siniscalchi \(1999, Footnote 5, p.193\)](#)).

¹⁶Where, in particular, the nature of the spaces in [Definition 3.2](#) is the main reason behind the need to introduce this construct. See also [Section 5](#) for various examples regarding the nature of the conditioning events, where, in particular, it is important to stress, as can be observed from [Section 5.3](#), that these conditioning events do not need to represent verifiable information (e.g., information sets reached in a dynamic game), but they should be considered simply a tool to capture counterfactual reasoning.

¹⁷Obviously, in absence of conditioning events, this definition boils down to the standard one of image measure as in [Aliprantis & Border \(2006, Chapter 13.12, p.483\)](#) or [Bogachev \(2007, Chapter 3.6, p.190\)](#).

Definition 2.4 (Image Measure with Conditioning Events). *Given a conditional measurable space $(M, \Sigma_M, \mathcal{C}_M)$, two product conditional measurable spaces $(M \times X, \Sigma_M \otimes \Sigma_X, \mathcal{C}_M)$ and $(M \times Y, \Sigma_M \otimes \Sigma_Y, \mathcal{C}_M)$ sharing \mathcal{C}_M , and a measurable function $f \in (M \times Y)^{M \times X}$, the image measure with conditioning events of f is the function*

$$\widehat{f} := (\widehat{f}_C)_{C \in \mathcal{C}_M} : \Delta^{\mathcal{C}_M}(M \times X) \rightarrow \Delta^{\mathcal{C}_M}(M \times Y),$$

defined as

$$\widehat{f}_C(\nu)(E) := \nu(f^{-1}(E)|C),$$

for every $C \in \mathcal{C}_M$, $\nu := (\nu(\cdot|C))_{C \in \mathcal{C}_M} \in \Delta^{\mathcal{C}_M}(M \times X)$, and $E \in \Sigma_M \otimes \Sigma_Y$.

In the following, it is understood that a given product conditional measurable space $(M \times X, \Sigma_M \otimes \Sigma_X, \mathcal{C}_M)$ induced by \mathcal{C}_M is endowed with

$$\Sigma_{\Delta^{\mathcal{C}_M}(M \times X)} := \sigma(\{ \gamma_C^p(E) \mid E \in \Sigma_M \otimes \Sigma_X, C \in \mathcal{C}_M, p \in [0, 1] \}) \quad (2.3)$$

naturally defined as an extension of [Definition 2.2](#) to $(M \times X)$.

2.2 Theory of Coalgebras

In what follows, it is assumed a minimal knowledge of certain notions from category theory:¹⁸ in particular, that of category (along with the notions of objects and morphisms of a category, and of isomorphism and isomorphic objects), (full) subcategory, (endo)functor (along with the definition of four basic functors, i.e., the identity functor, the constant functor, the power set endofunctor, and the projection functor), and terminal object in a category, where—for self-containment purposes—the related definitions can all be found in [Appendix A](#). Also, we employ the following—common in the field—notational conventions.

Categorical Notational Conventions. Given an arbitrary functor $\Phi : \mathbf{C} \rightarrow \mathbf{C}'$ with $\text{Ob}(\mathbf{C})$ denoting the objects of the category \mathbf{C} and $\mathbf{C}(A, A')$ denoting the morphisms between two objects $A, A' \in \text{Ob}(\mathbf{C})$,¹⁹ we write $\Phi(X)$ to capture how the functor acts on an arbitrary $X \in \text{Ob}(\mathbf{C})$, whereas we write Φf (without brackets) to capture how it acts on an arbitrary morphism $f \in \mathbf{C}(A, A')$. Also, given two arbitrary functors Φ and Φ' , we omit brackets (unless needed to avoid ambiguous expressions) to capture how the first applies to the second, i.e., we write $\Phi\Phi'$. Regarding morphisms, given two arbitrary morphisms f and g , we indifferently write “ fg ” or “ $f \circ g$ ” (of course, assuming that the latter expressions are well-defined). Finally, in commutative diagrams, we represent unique morphisms via *dashed* arrows.

As a matter of fact, this is everything we actually need for the next definition, which is going to prove to be crucial for our endeavour, i.e., that of a coalgebra.²⁰

Definition 2.5 (Coalgebra). *Given a category \mathbf{C} and an endofunctor Φ on \mathbf{C} , an Φ -coalgebra is a tuple $\mathfrak{C} = \langle S, \kappa \rangle$ where:*

- $S \in \text{Ob}(\mathbf{C})$ is its carrier;
- the morphism $\kappa : S \rightarrow \Phi(S)$ is its transition.²¹

¹⁸See [Leinster \(2014\)](#) for an introduction to the topic, [Mac Lane \(1998\)](#) for a comprehensive introduction, or [Borceux \(1994\)](#) for a more advanced treatment.

¹⁹See [Definition A.1](#) for the definition of category and the related notation as employed here.

²⁰See [Jacobs & Rutten \(1997\)](#) for an introductory article or [Jacobs \(2017\)](#) for a textbook on the topic.

²¹On the contrary, an Φ -algebra is a tuple $\mathfrak{A} := \langle S, \alpha \rangle$ with $\alpha : \Phi(S) \rightarrow S$ (see also [Mac Lane \(1998, Chapter VI.2\)](#)). The fact that the orientation of the morphism α is reversed with respect to the morphism κ is the reason behind the name “coalgebra”. Indeed, in category theory (and related fields), the prefix “co” is used when the direction of the morphisms in a given category \mathbf{C} is reversed: this amounts at working with the corresponding category \mathbf{C}^{op} , whose objects coincide with the objects of \mathbf{C} and such that $f^{\text{op}} \in \mathbf{C}^{\text{op}}(A', A)$ if $f \in \mathbf{C}(A, A')$. In particular, this has an immediate implication, which goes along the name of “Principle of Duality”, that can be informally stated as follows: every categorical definition, theorem, or proof has a dual, obtained by reversing all the morphisms (see [Leinster \(2014, Remark 1.1.10, p.16\)](#)).

Thus, a coalgebra is a structure that extracts information from a carrier S via the transition κ .²² In light of this, it is important to capture the way in which, given two coalgebras, the structure of the first is preserved when ‘moving’ to the second. The next definition achieves this goal.

Definition 2.6 (Coalgebra Morphism & Coalgebra Isomorphism). *Given a category \mathbf{C} , an endofunctor Φ on \mathbf{C} , and two Φ -coalgebras $\mathcal{C} := \langle S, \kappa \rangle$ and $\mathcal{C}' := \langle S', \kappa' \rangle$, an Φ -coalgebra morphism is a morphism $\mu : S \rightarrow S'$ such that the following diagram*

$$\begin{array}{ccc} S & \xrightarrow{\mu} & S' \\ \kappa \downarrow & & \downarrow \kappa' \\ \Phi(S) & \xrightarrow{\Phi\mu} & \Phi(S') \end{array}$$

commutes. If μ is an isomorphism, then it is a Φ -coalgebra isomorphism.

Notation 2 (Category of Coalgebras of an Endofunctor). Given a category \mathbf{C} and an endofunctor Φ on \mathbf{C} , we let $\mathbf{CoAlg}(\Phi)$ denote the category of Φ -coalgebras, whose objects are Φ -coalgebras and whose morphisms are Φ -coalgebra morphisms.

In this paper, we focus on the existence of a terminal object²³ in the category of Φ -coalgebra for a given endofunctor Φ on a category \mathbf{C} , i.e., the so-called terminal Φ -coalgebra.

Definition 2.7 (Terminal Coalgebra). *Given a category \mathbf{C} and an endofunctor Φ on \mathbf{C} , a terminal Φ -coalgebra $\bar{\mathcal{C}} := \langle \bar{S}, \bar{\kappa} \rangle$ is a terminal object in $\mathbf{CoAlg}(\Phi)$, i.e., it is a Φ -coalgebra such that for every Φ -coalgebra $\mathcal{C} := \langle S, \kappa \rangle$ there exists a unique Φ -coalgebra morphism $\bar{\mu} : S \rightarrow \bar{S}$ such that the following diagram*

$$\begin{array}{ccc} S & \xrightarrow{\bar{\mu}} & \bar{S} \\ \kappa \downarrow & & \downarrow \bar{\kappa} \\ \Phi(S) & \xrightarrow{\Phi\bar{\mu}} & \Phi(\bar{S}) \end{array}$$

commutes.

When a terminal coalgebra exists in $\mathbf{CoAlg}(\Phi)$, then this object is *essentially* unique, in light of the next well-known result from category theory, where we use the symbol “ \cong ” to capture the existence of an isomorphism between two objects in a category.²⁴

Lemma 1. *Given a category \mathbf{C} , terminal objects $\mathbf{1}, \mathbf{1}' \in \mathbf{Ob}(\mathbf{C})$ are isomorphic, i.e., $\mathbf{1} \cong \mathbf{1}'$.*

Whereas—as pointed out above—the focus of this work is on establishing the existence of a terminal coalgebra given an endofunctor on a category, it is *not* always the case that there exists a terminal coalgebra. Indeed, for example, given the category **Set** and the power set endofunctor \mathcal{P} ,²⁵ there does not exist a terminal \mathcal{P} -coalgebra, i.e., there exists no terminal \mathcal{P} -coalgebra $\bar{\mathcal{C}} := \langle \bar{S}, \bar{\kappa} \rangle$ such that $\bar{S} \cong \mathcal{P}(\bar{S})$, which is an immediate consequence of Cantor’s theorem. However, when the terminal coalgebra on a given endofunctor on a category does exist, it is possible to exploit an important result from category theory and the theory of coalgebras known as *Lambek’s lemma*,²⁶ that we recall next.

Lemma 2 (Lambek (1968, Lemma 2.2, p.153)). *If $\bar{\mathcal{C}} := \langle \bar{S}, \bar{\kappa} \rangle$ is a terminal Φ -coalgebra, then the transition $\bar{\kappa} : \bar{S} \rightarrow \Phi(\bar{S})$ is an isomorphism, i.e., $\bar{S} \cong \Phi(\bar{S})$.*

²²See Jacobs (2017, Chapter 1) for an overview of how coalgebras arise in different contexts.

²³See Definition A.7 for the definition of terminal object.

²⁴See Definition A.5 for the definition of isomorphism and Definition A.6 for the definition of isomorphic objects.

²⁵See Example A.1 for the definition of the category **Set** and Example A.3 for the definition of the power set endofunctor.

²⁶In Lambek (1968), there is no mentioning of the underlying coalgebraic nature of the result. See also Jacobs (2017, Chapter 2.3, pp.69–70) for a textbook presentation of the result.

Thus, regarding the interpretation of Lemma 2, by focusing on the representation of the result as establishing $\bar{S} \cong \Phi(\bar{S})$, it is important to observe that this amounts at finding a *fixed point* for a given endofunctor Φ .

3. TYPE STRUCTURES

In the following, we let I denote a finite set of *agents* (or individuals) and 0 stand for what is called “Nature”, with $0 \notin I$, from which we define $I_0 := I \cup \{0\}$: we typically use the symbol “ i ” for a representative element of I and “ j ” for a representative element of I_0 . Concerning these sets when used for indexing purposes, we adopt the following standard conventions:

- if I_0 is the index set, we let $X := \prod_{j \in I_0} X_j$ and $X_{-j} := \prod_{y \in I_0 \setminus \{j\}} X_y$;
- if I is the index set, we let $X := \prod_{i \in I} X_i$ and $X_{-i} := \prod_{y \in I \setminus \{i\}} X_y$.

Given a family of functions $(f_j)_{j \in I_0}$ of the form $f_j : X_j \rightarrow Y_j$ (with the same convention applied when the index set is I), the *induced function* $f : X \rightarrow Y$ is defined as

$$f((x_j)_{j \in I_0}) := (f_j(x_j))_{j \in I_0}.$$

Finally, we employ the following non-standard convention, that considerably lighten our notation: given an agent $i \in I$ and an $|I_0|$ -tuple of functions (f_0, f_i, f_{-i}) , we let $f_{\pm i} := (f_0, f_{-i})$.

3.1 Basic Definition

For our endeavour, first of all, we fix a conditional measurable space $(\Theta, \Sigma_\Theta, \mathcal{C}_\Theta)$ that captures the common domain of uncertainty of interest. We now provide a formal definition of what a type structure is on $(\Theta, \Sigma_\Theta, \mathcal{C}_\Theta)$.

Definition 3.1 (Type Structure). *A type structure on a conditional measurable space $(\Theta, \Sigma_\Theta, \mathcal{C}_\Theta)$ is a tuple*

$$\mathcal{T} := \langle (T_i, \beta_i)_{i \in I} \rangle$$

where, for every $i \in I$,

- T_i is a measurable space, called the type space²⁷ of agent i ;
- $\beta_i := (\beta_{i,C})_{C \in \mathcal{C}_\Theta} : T_i \rightarrow \Delta^{\mathcal{C}_\Theta}(\Theta \times T_{-i})$ is a measurable function, called the belief function of agent i .

Thus, given that $t := (t_i)_{i \in I}$, an element (θ, t) is called a *state of the world*, with $\Theta \times T$ called the set of states of the world or *state space*, while a $t_i \in T_i$ is called an *epistemic type* (henceforth, type) of agent $i \in I$, for every $i \in I$. In particular, for every $i \in I$ and type $t_i \in T_i$, the belief function $\beta_i(t_i)$ captures the belief of type t_i regarding $\Theta \times T_{-i}$, for every conditioning event $C \in \mathcal{C}_\Theta$, i.e., $\beta_{i,C}(t_i) \in [\Delta(\Theta \times T_{-i})]^{\{C\}} = \Delta(\Theta \times T_{-i})$, for every $C \in \mathcal{C}_\Theta$.

3.2 Relation Between Type Structures

Having formalized the notion of type structure, we now provide a formal definition of a morphism between two type structures on a conditional measurable space $(\Theta, \Sigma_\Theta, \mathcal{C}_\Theta)$ that preserves the properties of the spaces belonging to the two structures.

Definition 3.2 (Type Morphism & Type Isomorphism). *Given a conditional measurable space $(\Theta, \Sigma_\Theta, \mathcal{C}_\Theta)$, two type structures $\mathcal{T} := \langle (T_i, \beta_i)_{i \in I} \rangle$ and $\mathcal{T}' := \langle (T'_i, \beta'_i)_{i \in I} \rangle$ on the conditional measurable space $(\Theta, \Sigma_\Theta, \mathcal{C}_\Theta)$, the induced measurable function $\vartheta := (\vartheta_j)_{j \in I_0}$ with $\vartheta_0 : \Theta \rightarrow \Theta$ and $\vartheta_i : T_i \rightarrow T'_i$, for every $i \in I$, is a type morphism if*

²⁷In the literature, the word “space” is often used to refer both to a tuple of objects such as \mathcal{T} and the type spaces T_i , for every $i \in I$. Here, we distinguish these constructs by using the word “structure” for the tuple and the word “space” for the set of types.

1) $\vartheta_0 := \mathbf{id}_\Theta$;

2) for every $i \in I$, $\beta'_i \circ \vartheta_i = \widehat{\vartheta}_{\pm i} \circ \beta_i$, i.e., the diagram

$$\begin{array}{ccc} T_i & \xrightarrow{\vartheta_i} & T'_i \\ \beta_i \downarrow & & \downarrow \beta'_i \\ \Delta^{\mathcal{C}_\Theta}(\Theta \times T_{-i}) & \xrightarrow{\widehat{\vartheta}_{\pm i}} & \Delta^{\mathcal{C}_\Theta}(\Theta \times T'_{-i}) \end{array}$$

commutes.

If $(\vartheta_j)_{j \in I_0}$ is a measurable isomorphism, then the type morphism is a type isomorphism.

In particular, Condition (2) in [Definition 3.2](#) can be alternatively formalized by saying that

$$\beta'_{i,C}(\vartheta_i(t_i))(E) = \beta_{i,C}(t_i)(\vartheta_{\pm i}^{-1}(E))$$

for every $i \in I$, $t_i \in T_i$, $C \in \mathcal{C}_\Theta$, and $E \subseteq \Theta \times T'_{-i}$.

3.3 Infinite Hierarchies of Beliefs

Given a conditional measurable space $(\Theta, \Sigma_\Theta, \mathcal{C}_\Theta)$ and a type structure $\mathcal{T} := \langle (T_i, \beta_i)_{i \in I} \rangle$ appended to it, for every $i \in I$, we can unpack the information contained in a type $t_i \in T_i$. This can be achieved by means of infinite hierarchies of beliefs.

The idea behind the construction of infinite hierarchies of beliefs is to construct beliefs of increasing order, and, for every belief order, a corresponding family of conditioning events.

Definition 3.3 (The Hierarchical Space). For every $i \in I$, proceed with the following inductive construction, starting from $H_i^0 := \{\bullet\} = \mathcal{C}^0$, for every $m \in \mathbb{N} := \{0, 1, 2, \dots\}$:

$$\begin{array}{ccc} H_i^1 := \Delta^{\mathcal{C}_i^1}(\Theta), & \mathcal{C}_i^1 := \mathcal{C}_\Theta, \\ \vdots & \vdots \\ H_i^{m+1} := \Delta^{\mathcal{C}_i^m}(\Theta \times H_{-i}^m), & \mathcal{C}_i^{m+1} := \{ D \subseteq \Theta \times H_{-i}^m \mid \exists C \in \mathcal{C}^m : D = C \times H_{-i}^m \} = \mathcal{C}_\Theta, \\ \vdots & \vdots \end{array}$$

The hierarchical space of agent i is $H_i := \prod_{\ell=0}^{\infty} H_i^\ell$, with $H := \prod_{i \in I} H_i$ the hierarchical space.²⁸

We now introduce a family of functions, one for every agent $i \in I$, to extract information from the hierarchical space. Thus, in the following, we let $!_\Theta$ denote the—unique—function $!_\Theta : \Theta \rightarrow \{\bullet\}$.

Definition 3.4 (Hierarchy Functions). Given a type structure $\mathcal{T} := \langle (T_i, \beta_i)_{i \in I} \rangle$ and an arbitrary agent $i \in I$, the function

$$\chi_i^{\beta,m} := (\chi_{i,C}^{\beta,m})_{C \in \mathcal{C}_\Theta} : T_i \rightarrow H_i^m$$

is the m -th order hierarchy function of $i \in I$, inductively defined as follows, for every $m \in \mathbb{N}$, $C \in \mathcal{C}_\Theta$, and $t_i \in T_i$:

- ($m = 0$) Let $\chi_{i,C}^{\beta,0}$ be uniquely defined as $\chi_{i,C}^{\beta,0} : T_i \rightarrow \{\bullet\}$;
- ($m \geq 1$) Assume that $\chi_C^{\beta,m} := (\chi_{i,C}^{\beta,m})_{i \in I}$ has been defined and let $\chi_{i,C}^{\beta,m+1}$ be defined as

$$\begin{aligned} \chi_{i,C}^{\beta,m+1}(t_i) &:= \left(\chi_{i,C}^{\beta,0}(t_i), \beta_{i,C}(t_i) \circ ((!_\Theta)^{-1}, (\chi_{-i,C}^{\beta,m})^{-1}) \right), \\ &= \left(\chi_{i,C}^{\beta,0}(t_i), \beta_{i,C}(t_i) \left(\Theta \times (\chi_{-i,C}^{\beta,m})^{-1}(t_i) \right) \right), \end{aligned}$$

for every $C \in \mathcal{C}_\Theta$.

²⁸It is understood that in this definition, in the part that pertains the conditioning events, we exploit [Notation 1](#).

Thus, the hierarchy function of agent i is the function $\chi_i^\beta := (\chi_{i,C}^\beta)_{C \in \mathcal{C}_\Theta} : T_i \rightarrow H_i$ such that $\chi_{i,C}^{\beta,m} = \pi_m \circ \chi_{i,C}^\beta$, for every $m \in \mathbb{N}$ and $t_i \in T_i$. Finally, the hierarchy function is the unique induced function

$$\chi^\beta := (\chi_{i,C}^\beta)_{i \in I, C \in \mathcal{C}_\Theta} : T \rightarrow H,$$

where an element $\chi_C^\beta(t)$ is deemed the hierarchy description of $t \in T$ at C .

3.4 Large Type Structures

In light of what we introduced in the previous sections, we can now collect the definitions of large type structures that we employ in this paper.

The notion of terminality we use goes back to [Armbruster & Böge \(1979, Section 4, p.19\)](#)²⁹ and [Böge & Eisele \(1979, Section 2, p.196\)](#), that employ this terminology remaining true to the original categorical³⁰ formulation of the problem at hand.³¹

Definition 3.5 (Terminal Type Structure). A type structure $\overline{\mathcal{T}} := \langle \langle \overline{T}_i, \overline{\beta}_i \rangle_{i \in I} \rangle$ is terminal if for every type structure \mathcal{T} on $(\Theta, \Sigma_\Theta, \mathcal{C}_\Theta)$ there exists a unique type morphism from \mathcal{T} to $\overline{\mathcal{T}}$.

The next notion has been introduced in the literature as *completeness*³² in [Brandenburger \(2003\)](#).

Definition 3.6 (Belief-Complete Type Structure). A type structure $\overline{\mathcal{T}} := \langle \langle \overline{T}_i, \overline{\beta}_i \rangle_{i \in I} \rangle$ is belief-complete if the function $\overline{\beta}_i$ is surjective, for every $i \in I$.

The next definition, introduced in [Mertens & Zamir \(1985, Definition 2.4, p.6\)](#), formally captures the idea that, given a type structure $\mathcal{T} := \langle \langle T_i, \beta_i \rangle_{i \in I} \rangle$, the hierarchy function χ_i^β is injective, for every $i \in I$, i.e., for every $t_i, t'_i \in T_i$, if $t_i \neq t'_i$, then $\chi_i^\beta(t_i) \neq \chi_i^\beta(t'_i)$,³³ where, in particular, it should be observed how [Liu \(2009, Proposition 2, p.2123\)](#) establishes that the injectivity of the (induced) hierarchy function of a given type structure actually characterizes its non-redundancy.

Definition 3.7 (Non-redundancy). Given a type structure $\overline{\mathcal{T}} := \langle \langle \overline{T}_i, \overline{\beta}_i \rangle_{i \in I} \rangle$ and the corresponding (induced) hierarchy function $\chi^{\overline{\beta}} := (\chi_i^{\overline{\beta}})_{i \in I}$, $\overline{\mathcal{T}}$ is non-redundant if the σ -algebra $\sigma(\chi_i^{\overline{\beta}})$ on \overline{T}_i is separating, for every $i \in I$.³⁴

Thus, extending a terminology introduced in [Siniscalchi \(2008\)](#) and hinted in [Di Tillio \(2008, Section 3.4\)](#), we now define the so-called universal type structure.³⁵

Definition 3.8 (Universal Type Structure). A type structure $\overline{\mathcal{T}} := \langle \langle \overline{T}_i, \overline{\beta}_i \rangle_{i \in I} \rangle$ is universal if it is terminal, belief-complete, and non-redundant.

²⁹Given the general lack of availability of this paper, we report here what we consider the relevant part with respect to the point above. Thus, interestingly, given that in [Armbruster & Böge \(1979\)](#) an oracle system is essentially a type structure, in [Armbruster & Böge \(1979, Section 4, p.19\)](#), we find that “the canonical oracle system is a terminal object in the category of oracle systems and n -tuples of commuting maps”, where the authors use the expression “canonical oracle system” essentially to refer to the universal type structure built from infinite hierarchies of coherent beliefs (as a projective limit) and the expression “ n -tuples of commuting maps” to refer to an n -tuple of type isomorphisms, with $n \in \mathbb{N}$ being the cardinality of the set of agents. We are grateful to Michael Greinecker for having made available to us the original article.

³⁰The same terminology can be found in [Vassilakis \(1991\)](#), [Vassilakis \(1992\)](#), and [Pintér \(2010\)](#), that refer explicitly to a categorical reformulation of the problem at hand.

³¹It should be pointed out how two other notions of terminality can be found in the literature, namely, in [Siniscalchi \(2008, p.93\)](#) (built on the intuition in [Battigalli & Siniscalchi \(1999, Remark 2, p.201\)](#)) and [Friedenberg \(2010, Definition 2.6, p.114\)](#). We are grateful to Gabriel Ziegler for having raised our attention to this point.

³²With [Siniscalchi \(2008\)](#) adopting the same term. By employing this terminology, we follow [Meier \(2012\)](#). Regarding this notion, see [Section 6.2.2](#).

³³See in general the discussion in [Mertens & Zamir \(1985, Section 2, p.6\)](#).

³⁴This notion has been introduced in [Mertens & Zamir \(1985, Definition 2.4, p.6\)](#). See in general the discussion in [Mertens & Zamir \(1985, Section 2, p.6\)](#) to motivate it and [Liu \(2009\)](#).

³⁵See [Section 6.2.3](#) for a discussion of this notion by considering that an alternative notion of universality can be found in the literature, as in [Brandenburger & Keisler \(2006, Section 11.i\)](#) and [Friedenberg \(2010, Section 5.a\)](#).

4. UNIVERSALITY

The main goal of this paper is to show that, for every conditional measurable space $(\Theta, \Sigma_\Theta, \mathcal{C}_\Theta)$, there exists a type structure appended to it that is universal as in [Definition 3.8](#).

In this section we provide a proof of the following theorem, where, for the case of a measurable space in absence of conditioning events, terminality, uniqueness, and non-redundancy have been proved by [Heifetz & Samet \(1998\)](#), whereas the belief-completeness has been proved by [Meier \(2012\)](#), [Moss & Viglizzo \(2004\)](#), and [Viglizzo \(2005b\)](#).

Theorem 1. *For every conditional measurable space $(\Theta, \Sigma_\Theta, \mathcal{C}_\Theta)$, there exists a type structure $\mathcal{T}^* := \langle (T_i^*, \beta_i^*)_{i \in I} \rangle$ on $(\Theta, \Sigma_\Theta, \mathcal{C}_\Theta)$ such that:*

- 1) Terminality: \mathcal{T}^* is terminal;
- 2) Uniqueness: \mathcal{T}^* is unique up to measurable type isomorphism;
- 3) Belief-Completeness: \mathcal{T}^* is belief-complete;
- 4) Non-Redundancy: \mathcal{T}^* is non-redundant.

The type structure \mathcal{T}^ is the Universal Type Structure on $(\Theta, \Sigma_\Theta, \mathcal{C}_\Theta)$, which is unique up to measurable type isomorphism.*

To establish this result, we employ the following strategy. First of all, in [Section 4.1](#), we bypass the presence of interactive agents and we show that there is essentially a basic mathematical structure underlying the very notion of type structure (in presence of conditioning events), which is nothing more than a coalgebra for an appropriate endofunctor. As a result, we establish the existence of a terminal coalgebra for this endofunctor in [Proposition 2](#). Armed with this result, in [Section 4.2](#), we tackle the presence of interactive agents and we show that the main bulk of the endeavour has—essentially—already been established. Indeed, a minimal change in the definition of the appropriate functor is enough to establish [Proposition 3](#), which is a translation of [Theorem 1](#) in coalgebraic terms as set in [Proposition 2](#) (i.e., taking into account the presence of interactive agents). Finally, in [Section 4.3](#), we simply show that type structures are indeed coalgebras for the appropriate endofunctor defined in [Section 4.2](#), thus exploiting the obtained result to establish [Theorem 1](#) essentially as its immediate corollary.

Before embarking in proceeding along the lines of the path sketched above, it is important to point out that the sections that follow have been written in a modular fashion. As a result, the reader who does not want to delve into the details of the coalgebraic canonical construction along with its structural properties as described in details in [Section 4.1](#) can skip that section and move to [Section 4.2](#), where there is essentially a description of type structures as a certain kind of coalgebras, which is actually the very point established in [Section 4.3](#) which delivers the theorem above.

4.1 There (in the Coalgebraic Framework)...

The very first step of our endeavour is to introduce the category that is going to be the main building block behind our undertaking.

Definition 4.1 (Category Meas). *The category **Meas** is the category of measurable spaces, where*

- *the objects of **Meas** are measurable spaces;*
- *the morphisms of **Meas** are measurable functions.*

Now, given that we fix a conditional measurable space $(M, \Sigma_M, \mathcal{C}_M)$, where—of course—we have that $M \in \text{Ob}(\mathbf{Meas})$, the first two functors needed for our endeavour are the constant functor \mathbf{K}_M and the identity functor \mathbf{Id} .³⁶ As a matter of fact, they are the building blocks of the derived

³⁶See [Example A.3](#) in [Appendix A](#) for their definitions, where it should be observed how, from a typographical standpoint, we distinguish the identity *endofunctor* from the identity *morphism* via the usage in the former of an *uppercase* initial letter (i.e., “F”).

functor \mathbf{P}_M , which is the functor defined as the binary product of the constant functor \mathbf{K}_M and the identity functor \mathbf{Id} , i.e., $\mathbf{P}_M := \mathbf{K}_M \times \mathbf{Id}$, where

- for every $X \in \mathbf{Ob}(\mathbf{Meas})$, $\mathbf{P}_M(X) := M \times X$,
- and

$$\mathbf{P}_M f : M \times X \rightarrow M \times Y$$

such that $\mathbf{P}_M f := (\mathbf{id}_M, f)$, for every $f \in \mathbf{Meas}(X, Y)$.

However, the category \mathbf{Meas} is somewhat too ‘large’ to capture the peculiarities of product conditional measurable spaces and functions between them. The definition that follows introduces a (full) subcategory³⁷ of \mathbf{Meas} which—on the contrary—is perfectly suited for this objective.

Definition 4.2 (Subcategory \mathbf{Meas}_M). *Given a conditional measurable space $(M, \Sigma_M, \mathcal{C}_M)$, the full subcategory \mathbf{Meas}_M of \mathbf{Meas} consists of*

- product conditional measurable spaces sharing \mathcal{C}_M and
- the measurable functions between product conditional measurable spaces sharing \mathcal{C}_M .

Now, given a (conditional) measurable space $(M, \Sigma_M, \mathcal{C}_M)$, building on [Definition 2.4](#), we introduce the endofunctor $\Delta^{\mathcal{C}_M}$ on \mathbf{Meas}_M defined as

- for every $M \times X \in \mathbf{Ob}(\mathbf{Meas}_M)$, $\Delta^{\mathcal{C}_M}(M \times X) := \Delta^{\mathcal{C}_M}(M \times X)$,
- and

$$\Delta^{\mathcal{C}_M} f := \widehat{f} : \Delta^{\mathcal{C}_M}(M \times X) \rightarrow \Delta^{\mathcal{C}_M}(M \times Y)$$

as in [Definition 2.4](#), for every $f \in \mathbf{Meas}_M(M \times X, M \times Y)$.

That this is—indeed—an endofunctor³⁸ is immediate in light of the measurability of the image measure.

Definition 4.3 (Type Functor). *The type functor $\overline{\mathbf{T}}$ is the morphism defined as*

$$\overline{\mathbf{T}} := \Delta^{\mathcal{C}_M} \mathbf{P}_M = \Delta^{\mathcal{C}_M}(\mathbf{K}_M \times \mathbf{Id}),$$

such that

- for every $X \in \mathbf{Ob}(\mathbf{Meas})$, $\overline{\mathbf{T}}(X) := \Delta^{\mathcal{C}_M}(M \times X)$,
- and

$$\overline{\mathbf{T}} f : \Delta^{\mathcal{C}_M}(M \times X) \rightarrow \Delta^{\mathcal{C}_M}(M \times Y)$$

as in [Definition 2.4](#), for every $f \in \mathbf{Meas}(X, Y)$.

Given the setting above, we can now state a coalgebraic version of [Theorem 1](#), where the remainder of this section is devoted to establish this result.

Proposition 2. *Fix a conditional measurable space $(M, \Sigma_M, \mathcal{C}_M)$.*

- 1) Existence: *there exists a terminal $\overline{\mathbf{T}}$ -coalgebra $\mathfrak{Z} := \langle Z, \zeta \rangle$;*
- 2) Uniqueness: *the terminal $\overline{\mathbf{T}}$ -coalgebra $\mathfrak{Z} := \langle Z, \zeta \rangle$ is unique up to $\overline{\mathbf{T}}$ -coalgebra isomorphism;*
- 3) Isomorphic Transitions: *the transition ζ of the terminal $\overline{\mathbf{T}}$ -coalgebra $\mathfrak{Z} := \langle Z, \zeta \rangle$ is an isomorphism.*

In other words, in line with the interpretation provided in [Section 2.2](#) of terminal coalgebras as fixed point of a given functor, we want to establish the existence of an object $Z \in \mathbf{Ob}(\mathbf{CoAlg}(\overline{\mathbf{T}}))$ such that $Z \cong \overline{\mathbf{T}}(Z) \equiv \Delta^{\mathcal{C}_M}(M \times Z)$, i.e., such that the measurable spaces Z and $\Delta^{\mathcal{C}_M}(M \times Z)$ are isomorphic.³⁹

³⁷See [Definition A.2](#) for the definition of (full) subcategory.

³⁸That is, a morphism on the same category that satisfies Axioms (AC)–(AI) in [Definition A.3](#).

³⁹See [Moss \(2011, Section 2, p.30\)](#).

4.1.1 Obtaining the Terminal Carriers

The proof of [Proposition 2](#) proceeds along the following coalgebraic canonical construction, which builds on [Adámek & Koubek \(1979, Section A\)](#).⁴⁰ The crucial element to perform this canonical construction is to work with a category with a terminal object.⁴¹ It turns out that the category **Meas** has a terminal object, namely, the measurable space $(\mathbf{1}, \Sigma_{\mathbf{1}})$, which is a singleton set $\{\bullet\}$ endowed with the discrete (and trivial) σ -algebra, where—of course—working with $(\mathbf{1}, \Sigma_{\mathbf{1}})$ as a terminal object is without loss of generality, since all terminal objects in a given category are isomorphic from [Lemma 1](#).

Definition 4.4 (ω^{op} -Chain on the Endofunctor $\overline{\mathbf{T}}$). *An ω^{op} -chain⁴² on the endofunctor $\overline{\mathbf{T}}$ is a diagram of the form*

$$\mathbf{1} \leftarrow \overline{\mathbf{T}}^1(\mathbf{1}) \xleftarrow{\overline{\mathbf{T}}^1!} \overline{\mathbf{T}}^2(\mathbf{1}) \xleftarrow{\overline{\mathbf{T}}^2!} \dots, \quad (4.1)$$

with:

- concerning objects,

$$\begin{aligned} \overline{\mathbf{T}}^0(\mathbf{1}) &:= \mathbf{1}, \\ \overline{\mathbf{T}}^1(\mathbf{1}) &:= \Delta^{C_M}(M \times \mathbf{1}), \\ &\vdots \\ \overline{\mathbf{T}}^{n+1}(\mathbf{1}) &:= \overline{\mathbf{T}}(\overline{\mathbf{T}}^n(\mathbf{1})), \end{aligned}$$

where, for every $m \in \mathbb{N}$, the space $(\overline{\mathbf{T}}^m(\mathbf{1}), \Sigma_{\overline{\mathbf{T}}^m(\mathbf{1})})$ is a measurable space;

- concerning morphisms and their composition,

$$\begin{aligned} \overline{\mathbf{T}}^0! &:= !, \\ \overline{\mathbf{T}}^1! &: \Delta^{C_M}(M \times \Delta^{C_M}(M \times \mathbf{1})) \rightarrow \Delta^{C_M}(M \times \mathbf{1}), \\ &\vdots \\ \overline{\mathbf{T}}^{n+1}! &:= \overline{\mathbf{T}}\overline{\mathbf{T}}^n!. \end{aligned}$$

It should be observed how [Definition 4.4](#) is essentially in the spirit of the definition of *hierarchies* in [Heifetz & Samet \(1998, Section 5, p.335\)](#) and of [Definition 3.3](#), which is a recursive definition with a base step that is a singleton set (which in our context is exactly the terminal object $\mathbf{1}$).

Remark 4.1. *Since $\overline{\mathbf{T}}^1(\mathbf{1}) = \overline{\mathbf{T}}(\mathbf{1})$ is nonempty, the morphism $! : \overline{\mathbf{T}}(\mathbf{1}) \rightarrow \mathbf{1}$ is measurable and surjective, from which it follows the measurability and surjectivity of $\overline{\mathbf{T}}^m!$, for every $m \geq 1$.*

Definition 4.5 (**Projective Limit of the ω^{op} -chain on $\overline{\mathbf{T}}$**). *The projective limit of the ω^{op} -chain on the endofunctor $\overline{\mathbf{T}}$ is the set*

$$P := \left\{ (x_\ell) \in \prod_{\ell \in \mathbb{N}} \overline{\mathbf{T}}^\ell(\mathbf{1}) \mid \forall m \in \mathbb{N} \left(\overline{\mathbf{T}}^m!(x_{m+1}) = x_m \right) \right\}. \quad (4.2)$$

⁴⁰In [Adámek & Koubek \(1979\)](#), the result is established for algebras, but it applies by duality—as in [Footnote 21](#)—to coalgebras as well, as emphasized in [Barr \(1993\)](#) and [Adámek & Koubek \(1995, Example, p.60\)](#). The intuition for this construction has to be found—in light of the tight relation between posets and categories—in the construction of a least fixed point for directed complete partial orders or lattices, originally identified in [Scott \(1976, Section 5, p.552\)](#) in the context of λ -calculus (see also [Jacobs \(2017, Chapter 4.6, p.223\)](#)). [Smyth & Plotkin \(1982\)](#) builds a categorical framework to deal with the aforementioned constructions. It should be also pointed out that [Worrell \(2005\)](#) is a paper that establishes the result we are after by performing a different construction (see [Viglizzo \(2005b, Section 5, p.406\)](#) regarding this point).

⁴¹For example, this canonical construction cannot be used with bipointed sets (i.e., sets of the form $\{\perp\} \sqcup X \sqcup \{\top\}$ with X an arbitrary set), as employed in [Freyd \(2008\)](#) in his coalgebraic description of $[0, 1]$, since the corresponding category does *not* have a terminal object.

⁴²Alternatively called “ ω^{op} -sequence”, where the name of this construct comes from the fact that this is a sequence over the first infinite ordinal number ω .

In the following, we let $\pi_m : \prod_{\ell \in \mathbb{N}} \overline{\mathbf{T}}^\ell(\mathbf{1}) \rightarrow \overline{\mathbf{T}}^m(\mathbf{1})$ denote the projection operator on $\prod_{\ell \in \mathbb{N}} \overline{\mathbf{T}}^\ell(\mathbf{1})$ as canonically defined, for every $m \in \mathbb{N}$. Thus, we let P be endowed with the σ -algebra

$$\Sigma_P := \sigma\left(\left\{ \pi_m^{-1}(E) \mid m \in \mathbb{N}, E \in \Sigma_{\overline{\mathbf{T}}^m(\mathbf{1})} \right\}\right). \quad (4.3)$$

The next definition is in the spirit of the definition of the *description map* of Heifetz & Samet (1998, Section 5, p.336) and of Definition 3.4.

Definition 4.6 (Cone for $\overline{\mathbf{T}}$ -coalgebra \mathfrak{C}). Given a $\overline{\mathbf{T}}$ -coalgebra $\mathfrak{C} := \langle S, \kappa \rangle$, a cone for the $\overline{\mathbf{T}}$ -coalgebra \mathfrak{C} (or \mathfrak{C} -cone) is a recursively defined collection of morphisms

- $h_0^\kappa := ! : S \rightarrow \mathbf{1}$ and
- $h_m^\kappa : S \rightarrow \overline{\mathbf{T}}^m(\mathbf{1})$ such that $h_{m+1}^\kappa = \overline{\mathbf{T}}h_m^\kappa \circ \kappa$, for every $m \in \mathbb{N}$,

such that⁴³ there exists a unique morphism

$$h^\kappa : S \rightarrow P$$

such that $h_\ell^\kappa(s) = \pi_\ell \circ h^\kappa(s)$, for every $s \in S$, and $\ell \in \mathbb{N}$.

Remark 4.2. For every $m \in \mathbb{N}$, $h_m^\kappa = (\overline{\mathbf{T}}^m!) \circ h_{m+1}^\kappa$.⁴⁴

The following commutative diagram provides a perspicuous representation of the properties captured in Definition 4.6 and Remark 4.2

$$\begin{array}{ccc} S & \xrightarrow{\kappa} & \Delta^{\mathcal{C}_M}(M \times S) \\ h_m^\kappa \downarrow & \searrow h_{m+1}^\kappa & \downarrow \overline{\mathbf{T}}h_m^\kappa \\ \overline{\mathbf{T}}^m(\mathbf{1}) & \xleftarrow{\overline{\mathbf{T}}^m!} & \overline{\mathbf{T}}^{m+1}(\mathbf{1}) = \Delta^{\mathcal{C}_M}(M \times \overline{\mathbf{T}}^m(\mathbf{1})) \end{array}$$

for every $m \in \mathbb{N}$.

We now want to establish that, given two $\overline{\mathbf{T}}$ -coalgebras and a $\overline{\mathbf{T}}$ -coalgebra morphism between the two, cones ‘preserve’ the properties of these $\overline{\mathbf{T}}$ -coalgebras in a precise sense, as established in Heifetz & Samet (1998, Proposition 5.1, p.336) for what are called there “(hierarchies) descriptions”.

Lemma 3. Given two $\overline{\mathbf{T}}$ -coalgebras $\mathfrak{C} := \langle S, \kappa \rangle$ and $\mathfrak{C}' := \langle S', \kappa' \rangle$ and a $\overline{\mathbf{T}}$ -coalgebra morphism $\mu : S \rightarrow S'$, $h^\kappa = h^{\kappa'} \circ \mu$ is a $\overline{\mathbf{T}}$ -coalgebra morphism.

Building on Definition 4.6, we can now define what is going to turn out as the carrier of the terminal $\overline{\mathbf{T}}$ -coalgebra we are interested in, namely,

$$Z := \{ z \in P \mid \exists \mathfrak{C} := \langle S, \kappa \rangle \in \text{Ob}(\mathbf{CoAlg}(\overline{\mathbf{T}})) \exists s \in S : z = h^\kappa(s) \}, \quad (4.4)$$

which is endowed with the σ -algebra Σ_Z inherited from P .

4.1.2 Obtaining the Terminal Transition

To obtain the terminal transition, building on Equation (4.1), for $\mathbf{G} \in \{\overline{\mathbf{T}}, \mathbf{P}_M, \mathbf{K}_M, \mathbf{Id}\}$ we introduce the following auxiliary ω^{op} -chain:⁴⁵

$$\mathbf{G}(\mathbf{1}) \xleftarrow{\mathbf{G}!} \mathbf{G}\overline{\mathbf{T}}^1(\mathbf{1}) \xleftarrow{\mathbf{G}\overline{\mathbf{T}}^1!} \overline{\mathbf{T}}^2(\mathbf{1}) \xleftarrow{\mathbf{G}\overline{\mathbf{T}}^2!} \dots \quad (4.5)$$

⁴³The following is what is deemed in the literature a *universal property* (see Leinster (2014, Introduction) and Jacobs (2017, Chapter 1.2, p.7), or Aluffi (2009, Chapter I.5, p.33)).

⁴⁴See Viglizzo (2005b, Section 2.2, p.398) for the details, where the notation “ $(\overline{\mathbf{T}}^m!)$ ” refers to Definition 4.4.

⁴⁵The idea behind the set $\{\overline{\mathbf{T}}, \mathbf{P}_M, \mathbf{K}_M, \mathbf{Id}\}$ is to decompose a given functor in ‘more basic’ functors that should act as the *ingredients* upon which the functor is built as in Jacobs (2001) (see also Jacobs (2017, Chapter 6.5, p.379)), which is the path taken in Moss & Viglizzo (2004) and Viglizzo (2005b). See also Section 6.1.2.

As a result, for every $\mathbf{G} \in \{\overline{\mathbf{T}}, \mathbf{P}_M, \mathbf{K}_M, \mathbf{Id}\}$, we let $P_{\mathbf{G}}$ denote the resulting projective limit, with $\pi_n^{\mathbf{G}} : P_{\mathbf{G}} \rightarrow \mathbf{G}\overline{\mathbf{T}}^n(\mathbf{1})$ denoting the projection on $n \in \mathbb{N}$ as canonically defined. Now, for every $\overline{\mathbf{T}}$ -coalgebra $\mathfrak{C} := \langle S, \kappa \rangle$, there exists a \mathfrak{C} -cone with

$$\mathbf{G}h_n^{\kappa} : \mathbf{G}(S) \rightarrow \mathbf{G}\overline{\mathbf{T}}^n(\mathbf{1}),$$

for every $n \in \mathbb{N}$, and

$$h_{\mathbf{G}}^{\kappa} : \mathbf{G}(S) \rightarrow P_{\mathbf{G}}$$

such that $\pi_n^{\mathbf{G}} \circ h_{\mathbf{G}}^{\kappa} = \mathbf{G}h_n^{\kappa}$. Finally, we let $Z_{\mathbf{G}}$ be defined on $P_{\mathbf{G}}$ exactly like Z as in Equation (4.4) (of course, modulo the fact that we now work with Equation (4.5) and $P_{\mathbf{G}}$).

Thus, in particular, by spelling out the details:

- when $\mathbf{G} := \overline{\mathbf{T}}$, we have

$$\overline{\mathbf{T}}^1(\mathbf{1}) \xleftarrow{\overline{\mathbf{T}}^1!} \overline{\mathbf{T}}^2(\mathbf{1}) \xleftarrow{\overline{\mathbf{T}}^2!} \overline{\mathbf{T}}^3(\mathbf{1}) \xleftarrow{\overline{\mathbf{T}}^3!} \dots, \quad (4.6)$$

which corresponds to

$$\Delta^{\mathcal{C}_M}(M \times \mathbf{1}) \xleftarrow{\overline{\mathbf{T}}^1!} \Delta^{\mathcal{C}_M}(M \times (\Delta^{\mathcal{C}_M}(M \times \mathbf{1}))) \xleftarrow{\overline{\mathbf{T}}^2!} \Delta^{\mathcal{C}_M}(M \times \Delta^{\mathcal{C}_M}(M \times (\Delta^{\mathcal{C}_M}(M \times \mathbf{1})))) \xleftarrow{\overline{\mathbf{T}}^3!} \dots;$$

- when $\mathbf{G} := \mathbf{P}_M$, we have

$$\mathbf{P}_M(\mathbf{1}) \xleftarrow{\mathbf{P}_M!} \mathbf{P}_M\overline{\mathbf{T}}^1(\mathbf{1}) \xleftarrow{\mathbf{P}_M\overline{\mathbf{T}}^1!} \mathbf{P}_M\overline{\mathbf{T}}^2(\mathbf{1}) \xleftarrow{\mathbf{P}_M\overline{\mathbf{T}}^2!} \dots,$$

which corresponds to

$$M \times \mathbf{1} \xleftarrow{(\mathbf{id}_M, !)} M \times \overline{\mathbf{T}}^1(\mathbf{1}) \xleftarrow{(\mathbf{id}_M, \overline{\mathbf{T}}^1!)} M \times \overline{\mathbf{T}}^2(\mathbf{1}) \xleftarrow{(\mathbf{id}_M, \overline{\mathbf{T}}^2!)} \dots;$$

- when $\mathbf{G} := \mathbf{K}_M$, we have

$$\mathbf{K}_M \xleftarrow{\mathbf{K}_M!} \mathbf{K}_M\overline{\mathbf{T}}^1(\mathbf{1}) \xleftarrow{\mathbf{K}_M\overline{\mathbf{T}}^1!} \mathbf{K}_M\overline{\mathbf{T}}^2(\mathbf{1}) \xleftarrow{\mathbf{K}_M\overline{\mathbf{T}}^2!} \dots,$$

which corresponds to

$$M \xleftarrow{\mathbf{id}_M} M \xleftarrow{\mathbf{id}_M} M \xleftarrow{\mathbf{id}_M} \dots;$$

- when $\mathbf{G} := \mathbf{Id}$, we have

$$\mathbf{Id}(\mathbf{1}) \xleftarrow{\mathbf{Id}!} \mathbf{Id}\overline{\mathbf{T}}^1(\mathbf{1}) \xleftarrow{\mathbf{Id}\overline{\mathbf{T}}^1!} \mathbf{Id}\overline{\mathbf{T}}^2(\mathbf{1}) \xleftarrow{\mathbf{Id}\overline{\mathbf{T}}^2!} \dots,$$

which corresponds to

$$\mathbf{1} \xleftarrow{\overline{\mathbf{T}}^1!} \overline{\mathbf{T}}^1(\mathbf{1}) \xleftarrow{\overline{\mathbf{T}}^2!} \overline{\mathbf{T}}^2(\mathbf{1}) \xleftarrow{\overline{\mathbf{T}}^3!} \dots,$$

i.e., Equation (4.1) we started with, which implies that $Z_{\mathbf{Id}} = Z$ and $h_{\mathbf{Id}}^{\kappa} = h^{\kappa}$, for every $\overline{\mathbf{T}}$ -coalgebra $\mathfrak{C} := \langle S, \kappa \rangle$.

Remark 4.3. For every $k \in \mathbb{N}$, $\pi_k^{\mathbf{P}_M}(P_{\mathbf{P}_M}) = M \times \overline{\mathbf{T}}^k(\mathbf{1})$ and $\pi_k^{\mathbf{K}_M}(P_{\mathbf{K}_M}) = M^k$.

The importance of $Z_{\mathbf{G}}$, for every $\mathbf{G} \in \{\overline{\mathbf{T}}, \mathbf{P}_M, \mathbf{K}_M, \mathbf{Id}\}$, lies in the fact that we employ the following strategy to obtain the terminal transition:

- 1) we show the existence of a measurable morphism $\zeta_+ : Z \rightarrow Z_{\overline{\mathbf{T}}}$;
- 2) we show the existence of a measurable morphism $\overrightarrow{\zeta} : Z_{\overline{\mathbf{T}}} \rightarrow \overline{\mathbf{T}}(Z)$;
- 3) we let $\zeta : Z \rightarrow \overline{\mathbf{T}}(Z)$ be defined as $\zeta := \overrightarrow{\zeta} \circ \zeta_+$.

However, and rather crucially, Step (2) of the project relies on showing the existence of measurable morphisms $\zeta_{\mathbf{G}}$ and $\varphi_{\mathbf{G}}$, for every $\mathbf{G} \in \{\overline{\mathbf{T}}, \mathbf{P}_M, \mathbf{K}_M, \mathbf{Id}\}$. Essentially, the strategy revolves around the idea of capturing the very nature of the functor $\overline{\mathbf{T}}$ via morphisms $\zeta_{\mathbf{G}}$ (which capture the ‘nature’ of the functors upon which it is built) *and* morphisms $\varphi_{\mathbf{G}}$ (which encapsulate the order in which the functor ‘unpacks’ the functors that are its building blocks). In a more compact way, we want to obtain measurable spaces and measurable morphisms such that the following diagram

$$\begin{array}{ccc}
S & \xrightarrow{\kappa} & \overline{\mathbf{T}}(S) = \Delta^{c_M}(M \times S) \\
h^\kappa \downarrow & & \downarrow h_{\overline{\mathbf{T}}}^\kappa \\
Z & \xrightarrow{\zeta_+} Z_{\overline{\mathbf{T}}} & \xrightarrow{\overrightarrow{\zeta}} \overline{\mathbf{T}}(Z) = \Delta^{c_M}(M \times Z) \\
& \searrow \pi_{n+1} & \swarrow \overline{\mathbf{T}}\pi_n \\
& & \overline{\mathbf{T}}^{n+1}(\mathbf{1})
\end{array}
\quad (4.7)$$

$\overline{\mathbf{T}}h_n^\kappa$ (curved arrow from $\overline{\mathbf{T}}(S)$ to $\overline{\mathbf{T}}(Z)$)

commutes, with the understanding that $\overrightarrow{\zeta}$ is the result of the following diagram

$$\begin{array}{ccc}
Z_{\overline{\mathbf{T}}} = Z_{\Delta^{c_M} \mathbf{P}_M} & \xrightarrow{\zeta_{\Delta^{c_M} \mathbf{P}_M}} \Delta^{c_M}(Z_{\mathbf{P}_M}) & \xrightarrow{\Delta^{c_M} \zeta_{\mathbf{P}_M}} \Delta^{c_M}(Z_{\mathbf{K}_M} \times Z_{\mathbf{Id}}) \\
\zeta_+ \uparrow & & \downarrow \Delta^{c_M}(\zeta_{\mathbf{K}_M}, \zeta_{\mathbf{Id}}) \\
Z & \xrightarrow{\zeta} & \Delta^{c_M}(M \times Z)
\end{array}
\quad (4.8)$$

commuting, i.e., we want to define $\overrightarrow{\zeta}$ in such a way that Equation (4.8) commutes, where every $\zeta_{\mathbf{G}}$ for $\mathbf{G} \in \{\overline{\mathbf{T}}, \mathbf{P}_M, \mathbf{K}_M, \mathbf{Id}\}$ is defined in Appendix C and it is important to observe that the expressions “ $\Delta^{c_M}(Z_{\mathbf{P}_M})$ ” and “ $\Delta^{c_M}(Z_{\mathbf{K}_M} \times Z_{\mathbf{Id}})$ ” in Equation (4.8) are both well-defined in light of Remark 4.3. In particular, we want to define $\overrightarrow{\zeta}$ as

$$\begin{aligned}
\overrightarrow{\zeta} &:= \varphi_{\Delta^{c_M} \mathbf{P}_M} \\
&= \left(\Delta^{c_M} \varphi_{\mathbf{P}_M} \right) \zeta_{\Delta^{c_M} \mathbf{P}_M} \\
&= \left(\Delta^{c_M} \left((\zeta_{\mathbf{K}_M}, \zeta_{\mathbf{Id}}) \zeta_{\mathbf{P}_M} \right) \right) \zeta_{\Delta^{c_M} \mathbf{P}_M}
\end{aligned}
\quad (4.9)$$

where the idea behind Equation (4.9) is that, for example, in the passage from $\Delta^{c_M}(Z_{\mathbf{K}_M} \times Z_{\mathbf{Id}})$ to $\Delta^{c_M}(M \times Z)$, the functor Δ^{c_M} has already been employed in the previous step and as a result should not be written down, where the details can be found—once more—in Appendix C (in particular, in the proof of Lemma 5 below).⁴⁶

Thus, first of all, we establish Step (1) of the program above by proving the existence of a measurable morphism $\zeta_+ : Z \rightarrow Z_{\overline{\mathbf{T}}}$.

Lemma 4. *There exists a measurable morphism $\zeta_+ : Z \rightarrow Z_{\overline{\mathbf{T}}}$ such that $\zeta_+ \circ h^\kappa = h_{\overline{\mathbf{T}}}^\kappa \circ \kappa$, for every $\overline{\mathbf{T}}$ -coalgebra $\mathfrak{C} := \langle S, \kappa \rangle$.*

More succinctly, Lemma 4 can be captured by stating that the following diagram

$$\begin{array}{ccc}
S & \xrightarrow{\kappa} & \overline{\mathbf{T}}(S) \\
h^\kappa \downarrow & & \downarrow h_{\overline{\mathbf{T}}}^\kappa \\
Z & \xrightarrow{\zeta_+} & Z_{\overline{\mathbf{T}}}
\end{array}$$

commutes.

⁴⁶See Viglizzo (2005b, Example 2, p.405) for a similar reading of a commutative diagram in the same context.

Then, we establish Step (2) of the strategy above, with the understanding that all the—background—work whose intuition is sketched above regarding the existence of measurable morphisms $\zeta_{\mathbf{G}}$, for every $\mathbf{G} \in \{\bar{\mathbf{T}}, \mathbf{P}_M, \mathbf{K}_M, \mathbf{Id}\}$, can be found in [Appendix C](#).

Lemma 5. *There exists a measurable morphism*

$$\vec{\zeta} := (\vec{\zeta}_C)_{C_M} : Z_{\bar{\mathbf{T}}} \rightarrow \bar{\mathbf{T}}(Z) = \Delta^{C_M}(M \times Z)$$

such that:

- 1) $\vec{\zeta} \circ h_{\bar{\mathbf{T}}}^{\kappa} = \bar{\mathbf{T}}h^{\kappa}$, for every $\bar{\mathbf{T}}$ -coalgebra $\mathcal{C} := \langle S, \kappa \rangle$;
- 2) $\bar{\mathbf{T}}\pi_m \circ \vec{\zeta} = \pi_m^{\bar{\mathbf{T}}}$, for every $m \in \mathbb{N}$.

Once more, succinctly, [Lemma 5](#) amounts at establishing the commutativity of the following diagram

$$\begin{array}{ccc}
 & \Delta^{C_M}(M \times S) & \\
 h_{\bar{\mathbf{T}}}^{\kappa} \swarrow & & \searrow \bar{\mathbf{T}}h^{\kappa} \\
 Z_{\bar{\mathbf{T}}} & \xrightarrow{\vec{\zeta}} & \Delta^{C_M}(M \times Z) \\
 \pi_m^{\bar{\mathbf{T}}} \searrow & & \swarrow \bar{\mathbf{T}}\pi_m \\
 & \bar{\mathbf{T}}^{m+1}(\mathbf{1}) &
 \end{array} \tag{4.10}$$

where it should be recalled that $\bar{\mathbf{T}}^{m+1}(\mathbf{1}) = \Delta^{C_M}(M \times \bar{\mathbf{T}}^m(\mathbf{1}))$.

Finally, leveraging on [Lemma 4](#) and [Lemma 5](#), we now employ what we obtained with the previous lemmata by defining the *terminal transition* ζ as

$$\zeta := \vec{\zeta} \circ \zeta_+. \tag{4.11}$$

4.1.3 Establishing Terminality

We now collect what we established in [Section 4.1.1](#) and [Section 4.1.2](#) in a dedicated definition.

Definition 4.7 (The $\bar{\mathbf{T}}$ -Coalgebra \mathfrak{Z}). *The $\bar{\mathbf{T}}$ -Coalgebra*

$$\mathfrak{Z} := \langle Z, \zeta \rangle$$

is the $\bar{\mathbf{T}}$ -Coalgebra whose carrier Z is defined as in [Equation \(4.4\)](#) and whose transition ζ is defined as in [Equation \(4.11\)](#).

What now remains to do is to establish the terminality of \mathfrak{Z} . Thus, first of all we start from the following lemma, whose proof is immediate in light of the fact that the following diagram

$$\begin{array}{ccccc}
 S & \xrightarrow{\kappa} & \bar{\mathbf{T}}(S) & & \\
 h^{\kappa} \downarrow & & h_{\bar{\mathbf{T}}}^{\kappa} \downarrow & & \searrow \bar{\mathbf{T}}h^{\kappa} \\
 Z & \xrightarrow{\zeta_+} & Z_{\bar{\mathbf{T}}} & \xrightarrow{\vec{\zeta}} & \bar{\mathbf{T}}(Z) \\
 & \searrow \zeta & & \swarrow \zeta & \\
 & & & &
 \end{array}$$

commutes from [Lemma 4](#) and [Lemma 5](#), which is in the spirit of [Heifetz & Samet \(1998, Proposition 5.3, p.337\)](#).

Lemma 6 (Cone as Coalgebra Morphism). *For every $\bar{\mathbf{T}}$ -coalgebra $\mathcal{C} := \langle S, \kappa \rangle$, h^{κ} is a $\bar{\mathbf{T}}$ -coalgebra morphism.*

Notation 3 (Cone for $\overline{\mathbf{T}}$ -Coalgebra \mathfrak{Z}). We let h denote the cone of $\mathfrak{Z} := \langle Z, \zeta \rangle$.

We now establish the fact that the cone h is nothing more than the identity function of Z , which is analogous to Heifetz & Samet (1998, Lemma 5.4, p.338).

Lemma 7. *Given the $\overline{\mathbf{T}}$ -coalgebra \mathfrak{Z} , $h = \text{id}_Z$.*

We can now state the terminality of \mathfrak{Z} in the following lemma, which is analogous to Heifetz & Samet (1998, Theorem 5.5, p.338).

Lemma 8. *The $\overline{\mathbf{T}}$ -coalgebra $\mathfrak{Z} := \langle Z, \zeta \rangle$ is a terminal $\overline{\mathbf{T}}$ -coalgebra.*

Finally, we provide the proof of Proposition 2.

Proof of Proposition 2. We establish the points in the order in which they are presented in the statement of the result.

- 1) *Existence:* This is an immediate consequence of Lemma 8.
- 2) *Uniqueness:* The uniqueness (in this case up to $\overline{\mathbf{T}}$ -coalgebra isomorphism) of a terminal object comes from Lemma 1.
- 3) *Isomorphic Transitions:* The fact that ζ is an isomorphism is an immediate consequence of Lambek's lemma, i.e., Lemma 2.

Thus, what is written above establishes the result. ■

4.2 ... In the Dale (between the Two Worlds)...

In the previous section, we established the existence of a terminal $\overline{\mathbf{T}}$ -coalgebra in the category **Meas**. However, we do not have to lose sight of the fact that what we are after is actually a proof of Theorem 1, which is a theorem regarding the existence of a type structure, i.e., an object that is intrinsically related to the presence of *multiple* interactive agents. Thus, quite simply, **Meas** is not rich enough for our purposes. Indeed, we need to work in a theoretical framework that captures the presence of multiple interactive agents. Thus, the definition that follows accomplishes exactly this.

Definition 4.8 (Category \mathbf{Meas}^I). *The category \mathbf{Meas}^I is the category of measurable spaces, where*

- *the objects of \mathbf{Meas}^I are an $|I|$ -fold product of measurable spaces;*
- *the morphisms of \mathbf{Meas}^I are an $|I|$ -fold product of measurable functions.*

Hence, in order to get closer to the original problem addressed for type structures built on a common domain of uncertainty $(\Theta, \Sigma_\Theta, \mathcal{C}_\Theta)$, from now on we focus on a given conditional measurable space $(\Theta, \Sigma_\Theta, \mathcal{C}_\Theta)$. Also, with an innocuous abuse of notation, we employ the same symbols we used in the previous section (modulo usage of Θ instead of M), with the caveat that—with a minimal change in comparison to the previous section—now we let $\mathbf{P}_\Theta := \mathbf{K}_\Theta \times \mathbf{Proj}_{-i}$, i.e., \mathbf{P}_Θ is defined by using the constant functor \mathbf{K}_Θ on Θ and the projection functor \mathbf{Proj} indexed by I .⁴⁷ Thus, given a conditional measurable space $(\Theta, \Sigma_\Theta, \mathcal{C}_\Theta)$, we let the morphism

$$\mathbf{T} : \mathbf{Meas}^I \rightarrow \mathbf{Meas}^I$$

such that $\mathbf{T} := \Delta^{\mathcal{C}_\Theta} \mathbf{P}_\Theta$, i.e.,

$$\mathbf{T} := \Delta^{\mathcal{C}_\Theta} \mathbf{P}_\Theta = \Delta^{\mathcal{C}_\Theta} (\mathbf{K}_\Theta \times \mathbf{Proj}_{-i}),$$

be defined as,

⁴⁷See Example A.2 for its definition.

- for every $(T_i)_{i \in I} \in \text{Ob}(\mathbf{Meas}^I)$,

$$\mathbf{T}((T_i)_{i \in I}) := \left(\Delta^{\mathcal{C}_\Theta}(\Theta \times T_{-i}) \right)_{i \in I},$$

- and

$$\mathbf{T}\mu := \left(\widehat{(\mathbf{id}_\Theta, \mu_{-i})} : \Delta^{\mathcal{C}_\Theta}(\Theta \times T_{-i}) \rightarrow \Delta^{\mathcal{C}_\Theta}(\Theta \times T'_{-i}) \right)_{i \in I} \quad (4.12)$$

for every $\mu := (\mu_i)_{i \in I} \in \mathbf{Meas}^I((T_i)_{i \in I}, (T'_i)_{i \in I})$,

where it is immediate to establish that this morphism is indeed a functor.

Notation 4. We let $\mathbf{H}_i := \mathbf{Proj}_i \mathbf{T}$, i.e., $\mathbf{H}_i(T_i) := \mathbf{Proj}_i \mathbf{T}((T_i)_{i \in I})$ and $\mathbf{H}_i \mu_i := \mathbf{Proj}_i \mathbf{T}((\mu_i)_{i \in I})$.

It has to be observed that Equation (4.12) is well-defined in light of the definition of \mathbf{T} , which implies that, given a certain profile $T := (T_i)_{i \in I}$, the application of the functor \mathbf{P}_Θ leads to $\Theta \times T$, which—in turn—is related to the fact that, for every $i \in I$ with $\mu_i : T_i \rightarrow T'_i$, $\mathbf{H}_i \mu_i = \widehat{(\mathbf{id}_\Theta, \mu_{-i})}$ is well-defined, because the image measure $\widehat{(\mathbf{id}_\Theta, \mu_{-i})}$ is defined on the induced function $(\mathbf{id}_\Theta, \mu_{-i}) : \Theta \times T_{-i} \rightarrow \Theta \times T'_{-i}$ which is a product conditional measurable space as needed for Definition 2.4.

Definition 4.9 (T-Coalgebra). A \mathbf{T} -coalgebra is a tuple

$$\mathfrak{C} := \langle (T_i, \beta_i)_{i \in I} \rangle$$

where T_i is the i -carrier of the \mathbf{T} -coalgebra with transition $\beta_i : T_i \rightarrow \Delta^{\mathcal{C}_\Theta}(\Theta \times T_{-i})$, for every $i \in I$.

We can now reformulate Proposition 2 for the present setting where the focus is on the category \mathbf{Meas}^I .

Proposition 3. Fix a conditional measurable space $(\Theta, \Sigma_\Theta, \mathcal{C}_\Theta)$.

- 1) Existence: there exists a terminal \mathbf{T} -coalgebra $\mathfrak{Z} := \langle (Z_i, \zeta_i)_{i \in I} \rangle$;
- 2) Uniqueness: the terminal \mathbf{T} -coalgebra $\mathfrak{Z} := \langle (Z_i, \zeta_i)_{i \in I} \rangle$ is unique up to $\overline{\mathbf{T}}$ -coalgebra isomorphism;
- 3) Isomorphic Transitions: the transition ζ_i of the terminal \mathbf{T} -coalgebra $\mathfrak{Z} := \langle (Z_i, \zeta_i)_{i \in I} \rangle$ is an isomorphism, for every $i \in I$.

Whereas we do not give the details of the proof, which essentially proceeds along the lines of that of Proposition 2 with the caveat that a considerably heavier notation is needed to deal with the presence of multiple agents, we sketch the path that should be taken by proceeding along the lines of what has been done in Section 4.1. Thus, in the following, everything starts from the—by now classical—canonical construction, with the understanding that $\mathbf{T}(\mathbf{1})$ (and the like) is used to denote the application of the endofunctor $\overline{\mathbf{T}}$ to an $(|I| - 1)$ -tuple of $\mathbf{1}$ (endowed with the discrete σ -algebra), whereas $\mathbf{T}!$ (and the like) is used to denote the application of the endofunctor \mathbf{T} to an $(|I| - 1)$ -tuple of morphisms !.

It is useful to ‘unpack’ the information contained in Definition 4.4 by showing how the ω^{op} -chains look for a T_i , with $i \in I$ arbitrary. Hence, we have

$$\mathbf{1} \leftarrow \text{---}^! \text{---} \mathbf{H}_i^1(\mathbf{1}) \xleftarrow{\mathbf{H}_i^1!} \mathbf{H}_i^2(\mathbf{1}) \xleftarrow{\mathbf{H}_i^2!} \dots,$$

which, by spelling out the details, amounts at

$$\mathbf{1} \leftarrow \text{---}^! \text{---} \Delta^{\mathcal{C}_\Theta} \left(\Theta \times \prod_{j \in I \setminus \{i\}} \mathbf{1} \right) \xleftarrow{\mathbf{H}_i^1!} \Delta^{\mathcal{C}_\Theta} \left(\Theta \times \prod_{j \in I \setminus \{i\}} \left(\Delta^{\mathcal{C}_\Theta}(\Theta \times \prod_{j \in I \setminus \{i\}} \mathbf{1}) \right) \right) \xleftarrow{\mathbf{H}_i^2!} \dots,$$

in light of the fact that $\prod_{j \in I \setminus \{i\}} \mathbf{1} = \mathbf{1}^{|I|-1}$. Once more, it is immediate to observe by inspection that this is in the spirit of the construction of [Heifetz & Samet \(1998, Section 5, p.335\)](#) in presence of conditioning events.

Getting into the details of the profile $P := (P_i)_{i \in I}$ and decomposing the information contained in [Equation \(4.2\)](#), we have

$$P_i := \left\{ (x_\ell) \in \prod_{\ell \in \mathbb{N}} \mathbf{H}_i^\ell(\mathbf{1}) \mid \forall m \in \mathbb{N} \left(\mathbf{H}_i^{m!}(x_{m+1}) = x_m \right) \right\}.$$

Now, given a \mathbf{T} -coalgebra $\mathfrak{C} := \langle (T_i, \beta_i)_{i \in I} \rangle$, we define the notion of *cone* over that \mathbf{T} -coalgebra, whose role is to map the information ‘implicitly’ contained in the carrier $(T_i)_{i \in I}$ to $(P_i)_{i \in I}$ as defined above. Thus, given a \mathbf{T} -coalgebra $\mathfrak{C} := \langle (T_i, \beta_i)_{i \in I} \rangle$, a *cone* for the \mathbf{T} -coalgebra \mathfrak{C} is a recursively defined collection of morphisms, for every $i \in I$ and $m \in \mathbb{N}$,

- $h_{i,0}^\beta := ! : T_i \rightarrow \mathbf{1}$,
- $h_{i,m}^\beta : T_i \rightarrow \mathbf{H}_i^m(\mathbf{1})$ such that $h_{i,m+1}^\beta = \mathbf{H}_i h_{i,m}^\beta \circ \beta_i$,

such that there exists a unique morphism

$$h_i^\beta : T_i \rightarrow P_i$$

such that $h_{i,\ell}^\beta(t_j) = \pi_\ell \circ h_i^\beta(t_i)$, for every $i \in I$, $t_i \in T_i$, and $\ell \in \mathbb{N}$. Finally, building on [Definition 4.6](#), we can now define, for every $i \in I$, the carrier of the terminal \mathbf{T} -coalgebra we are interested in, namely,

$$Z_i := \left\{ z_i \in P_i \mid \exists \mathfrak{C} := \langle (T_i, \beta_i)_{i \in I} \rangle \in \text{Ob}(\mathbf{CoAlg}(\mathbf{T})) \exists t_i \in T_i : z_i = h_i^\beta(t_i) \right\},$$

which is endowed with the σ -algebra Σ_{Z_i} inherited from P_i , with $Z := (Z_i)_{i \in I}$.

Having obtained the structure of $Z := (Z_i)_{i \in I}$, constructing the terminal transition $\zeta := (\zeta_i)_{i \in I}$ and establishing the terminality of the corresponding \mathbf{T} -coalgebra proceeds along the same path taken in [Section 4.1.2](#) and [Section 4.1.3](#). In particular, given that we let $h := (h_i)_{i \in I}$ denote the cone of the terminal \mathbf{T} -coalgebra $\mathfrak{Z} := \langle (Z_i, \zeta_i)_{i \in I} \rangle$, we can establish the following crucial—and obviously related—properties of h_i , for every $i \in I$.

Remark 4.4. *For every $i \in I$, the σ -algebra on Z_i is separative, with $h_i = \mathbf{id}_{Z_i}$ obviously injective.*

4.3 ... and Back Again (to Type Structures)

We can now go back to our original problem, namely, the proof of [Theorem 1](#). In light of [Section 4.2](#), it should be observed how the notation we use in this section for the elements of a type structures and those of a coalgebra are actually the same. Thus, the proof of the next result is immediate by inspection. It has to be observed that this also—implicitly—establishes that cones (as in [Section 4.2](#)) are hierarchy functions as in [Definition 3.4](#).

Lemma 9 (Type Structures as Coalgebras). *Given a conditional measurable space $(\Theta, \Sigma_\Theta, \mathcal{C}_\Theta)$, a type structure $\mathcal{T} := \langle (T_i, \beta_i)_{i \in I} \rangle$ is a \mathbf{T} -coalgebra $\mathfrak{C} := \langle (T_i, \beta_i)_{i \in I} \rangle$.*

Crucially, we have to establish that type morphisms and coalgebra morphisms are actually the same, which we achieve next.

Lemma 10 (Type Morphisms as Coalgebra Morphisms). *Given a conditional measurable space $(\Theta, \Sigma_\Theta, \mathcal{C}_\Theta)$ and two arbitrary type structures $\mathcal{T} := \langle (T_i, \beta_i)_{i \in I} \rangle$, and $\mathcal{T}' := \langle (T'_i, \beta'_i)_{i \in I} \rangle$ on $(\Theta, \Sigma_\Theta, \mathcal{C}_\Theta)$ sharing \mathcal{C}_Θ , a type morphism $\vartheta := (\vartheta_j)_{j \in I_0}$ is a \mathbf{T} -coalgebra morphism.*

In light of what we established in [Section 4.1](#) and [Section 4.2](#) and the lemmata above, we can finally provide the proof of [Theorem 1](#).

Proof of Theorem 1. From Proposition 3, we have these immediate implications concerning Theorem 1:

- 1) *Terminality:* From Point (1) of Proposition 3, Lemma 9, and Lemma 10, there exists a terminal type structure

$$\mathcal{T}^* := \langle (T_i^*, \beta_i^*)_{i \in I} \rangle$$

on $(\Theta, \Sigma_\Theta, \mathcal{C}_\Theta)$;

- 2) *Uniqueness:* From Point (2) of Proposition 3, this terminal type structure \mathcal{T}^* is the unique up to measurable type isomorphism terminal type structure on $(\Theta, \Sigma_\Theta, \mathcal{C}_\Theta)$;

- 3) *Belief-Completeness:* From Point (3) of Proposition 3, the terminal type structure \mathcal{T}^* is belief-complete, i.e., the belief function β_i^* is surjective, for every $i \in I$.

- 4) *Non-Redundancy:* From Lemma 9 and Remark 4.4, \mathcal{T}^* is non-redundant by construction.

Being terminal, belief-complete, and non-redundant, \mathcal{T}^* is the Universal Type Structure on $(\Theta, \Sigma_\Theta, \mathcal{C}_\Theta)$ as in Definition 3.8, which, in light of its uniqueness up to measurable type isomorphism, establishes the result. \blacksquare

5. APPLICATIONS

On general and conceptual grounds, as hinted in Section 1.1, our construction can be used as a ‘foundation’ for the construction obtained in Battigalli & Siniscalchi (1999, Section 2). In order to see this point, all the relevant information has already been provided in Footnote 15, whose details we now spell out. Thus, starting from a σ -algebra over $\Delta^{\mathcal{C}_M}(M)$ as in Definition 2.2, we do provide a ‘conceptual’ foundation in terms of what a hypothetical individual could consider as a relevant event. And that is the crucial point: indeed, when we start from topological assumptions, as in Battigalli & Siniscalchi (1999), with $\Delta^{\mathcal{C}_M}(M)$ endowed with the topology of weak convergence, it turns out from Kechris (1995, Theorem 17.24, p.112) opportunely adapted to the presence of conditioning events that the two σ -algebras are one and the same. Thus, the events a hypothetical individual can conceive are ‘the same’ and—in a sense—we do not lose anything by working with topological assumptions in terms of *expressibility*.⁴⁸ It is actually in this spirit the usage of the present construction in Meier & Perea (2023)⁴⁹ for the epistemic characterization of their solution concept called “Forward and Backward Rationalizability”. And, once more, as already pointed out in Section 1.1, it is in this spirit that the present construction can be used for endeavours along the lines of Bergemann & Morris (2005), Dekel et al. (2007), or Battigalli et al. (2011) when working with conditioning events (similar, for example, to Penta (2015) and Müller (2016)).

However, crucially, as a matter of fact, we do lose something by working with topological assumptions when in presence of conditioning events, since the present work actually generalizes the one in Battigalli & Siniscalchi (1999, Section 2), where there is the need to assume an at most countable family of clopen (i.e., closed and open) conditioning events contrary to the present setting where the conditioning events are unrestricted.⁵⁰ Thus, in the next sections, we provide examples of how our construction can prove to be useful in various game-theoretical contexts by exploiting the two relaxations of the assumptions in Battigalli & Siniscalchi (1999, Section 2) we introduce, namely:

- lack of topological assumptions regarding the conditioning events;
- lack of cardinality assumptions regarding the conditioning events.

Additionally, we also show how the notion of conditioning event we employ proves to be amenable to various interpretations, a point which turns out to be useful—for example—for the study of environments in presence of lack of topological assumptions regarding a parameter space and ‘unverifiable’ conditioning events.

⁴⁸Regarding this notion, see for example Battigalli et al. (2011, Section 1.3).

⁴⁹With the caveat that the authors employ the term “universal” to refer to what we call here “terminal” (see Meier & Perea (2023, Section 4.1, p.20)).

⁵⁰See Section 6.3.3 for the details behind the need for these assumptions in Battigalli & Siniscalchi (1999) and our possibility of dropping them altogether.

5.1 Topology-Free Conditioning Events

A dynamic game with perfect recall and possibly simultaneous moves in its extensive form representation⁵¹ is a tuple

$$\Gamma := \langle I, (\Theta_i, A_i)_{i \in I}, X, Z, (H_i, S_i, u_i)_{i \in I} \rangle \quad (5.1)$$

where, given that the notation employed here has to be taken as standalone with respect to the rest of the paper, I now denotes the set of players, and, for every $i \in I$, we let Θ_i denote the set of *payoff types* of player i and A_i her set of *actions*. The set X is the set of *histories*, where a history x is either the empty sequence $\langle \emptyset \rangle$ (i.e., the initial history), or it is a sequence (a^1, \dots, a^K) , where $a^k := (a_i^k)_{i \in I}$ with $a_i^k \in A_i$ for every $i \in I$ and for every $1 \leq k \leq K$. We let $A(x) := \prod_{i \in I} A_i(x)$ denote the set of actions available to the players at history x , with player i *active* at history x if $|A_i(x)| \geq 2$. The set $Z \subseteq X$ is the set of *terminal* histories. Given that H_i denotes the set of *information sets* of player i with $H := \bigcup_{i \in I} H_i$ and that we extend to information sets the notation introduced above of actions available at a history, a *strategy* of player i is a function $s_i : H_i \rightarrow \bigcup_{h \in H_i} A_i(h)$ such that $s_i(h) \in A_i(h)$, for every $h \in H_i$. We let S_i denote the set of strategies of player i , with $S_{-i} := \prod_{j \in I \setminus \{i\}} S_j$ and $S := \prod_{j \in I} S_j$. Also, we let $S(h)$ denote the set of strategy profiles that allow information set $h \in H$, with $S_i(h) := \pi_{S_i} S(h)$ and $S_{-i}(h) := \pi_{S_{-i}} S(h)$. Finally, given that $\Xi_i := \Theta_i \times S_i$, with Ξ_{-i} , Ξ , and $\Xi_{-i}(h)$ accordingly defined as above, we let $u_i \in \mathfrak{R}^\Xi$ be a Bernoulli (unique up to positive affine transformation) utility function.

Thus, given the setting described above, as Battigalli & Tebaldi (2019, Section 5.6, p.758) point out, working with clopen conditioning events in the spirit of Battigalli & Siniscalchi (1999, Section 2) can be restrictive in certain applications. In particular, given a player $i \in I$ and assuming that Θ_i is uncountable and that player i obtains a signal regarding $\theta_{-i} \in \Theta_{-i}$, even if the signal is discrete, it is not necessarily the case that $\Xi_{-i}(h)$ is clopen for a given information set $h \in H_i$.⁵²

Now, it is important to observe that our construction covers the game-theoretical setting described above by freeing it from the necessity of working only with clopen conditioning events. Indeed, given that we let (Ξ, Σ_Ξ) be the relevant measurable space of interest, in the present context, we identify a player i 's set of conditioning events (topologically unrestricted) as the collection of sets $\Xi_{-i}(h)$, for every $h \in H_i$, i.e.,

$$\mathcal{C}_i := \{ \Xi_{-i}(h) \mid h \in H_i \},$$

with $\mathcal{C}_\Xi := \prod_{i \in I} \mathcal{C}_i$. Thus, building the universal type structure on the conditional measurable space $(\Xi, \Sigma_\Xi, \mathcal{C}_\Xi)$, we can capture a situation where a player i has beliefs over her own strategies and payoff types for every possible information set in the game, including those where she is not active. Now, if we additionally want to work with a type structure where every player i has beliefs only over *her* domain of uncertainty Ξ_{-i} and these beliefs arise only at those information set where she is active, i.e., those information sets belonging to H_i , we simply have to ‘carve out’ from the previously obtained universal type structure the corresponding belief-closed type structure where the only conditioning events are those in \mathcal{C}_i and the only relevant domain of uncertainty is given by Ξ_{-i} , for every i , where a *belief-closed* type structure⁵³ of a given type structure $\mathcal{T} := \langle (T_i, \beta_i)_{i \in I} \rangle$ is a type structure $\tilde{\mathcal{T}} := \langle (\tilde{T}_i, \tilde{\beta}_i)_{i \in I} \rangle$ such that $\text{supp } \tilde{\beta}_{i,C}(\tilde{t}_i) \subseteq \Xi_{-i} \times \tilde{T}_{-i}$, for every $i \in I$, $C \in \mathcal{C}_i$, and $\tilde{t}_i \in \tilde{T}_i$ with $\tilde{T}_i \subseteq T_i$. As a result, we would obtain, as mentioned above, a belief-closed type structure of the original one, which—nonetheless—would be itself universal in its own rights given the imposed restrictions.

5.2 Uncountability of the Conditioning Events

The notion of *Rationality and Common Strong Belief in Rationality* (henceforth, RCSBR) introduced in Battigalli & Siniscalchi (2002, Section 4.2, p.372) is the epistemic solution concept that

⁵¹See Osborne & Rubinstein (1994, Definition 200.1, Chapter 11.1.2) for a similar definition (under a different name), with the caveat that the latter does not explicitly allow for simultaneous moves (see Osborne & Rubinstein (1994, Definition 200.1, Chapter 6.3.2) for the corresponding extension).

⁵²See the similar discussion in De Vito (2023, Section 6).

⁵³The following notion is a translation in the framework of type structures of Mertens & Zamir (1985, Definition 2.15, p.12). Battigalli et al. (Work in Progress) refer to these constructs as *self-evident events*.

characterizes *Strong Rationalizability*⁵⁴ of Pearce (1984, Definition 9, p.1042) and Battigalli (1997, Definition 2, p.46). Recently, two papers have profitably employed RCSBR and Strong Rationalizability in applications: in particular, in Friedenberg (2019), RCSBR is used to study behavior in bargaining, while Strong Rationalizability is employed in Guo & Shmaya (2021, Proposition 2, p.495) to ‘refine’ the results obtained via Bayes Nash Equilibrium in the study of the behavior of a platform that provides probabilistic forecasts to a customer in presence of costly miscalibration. Interestingly, both papers point out, respectively, in Friedenberg (2019, Remark A.1, pp. 1855–1856) and Guo & Shmaya (2021, Section 4.2, p.494), a technical limitation of RCSBR and Strong Rationalizability, namely that both solution concepts are defined for an at most countable collection of conditioning events (i.e., information sets). Thus, with respect to this point, our construction can be the building block for a definition of RCSBR based on possibly uncountably many conditioning events, which, given an appropriate definition of Strong Rationalizability for possibly uncountably many conditioning events, could actually epistemically characterize the aforementioned new notion of Strong Rationalizability exploiting standard arguments in epistemic game theory based on the belief-completeness of the type structure under scrutiny.

5.3 Nature of the Conditioning Events

In both Section 5.1 and Section 5.2, we consider conditioning events as information sets of a dynamic game. However, as natural as this is, interestingly, information sets are *not* the only way in which we can interpret our *abstract* conditioning events. Thus, with respect to this point, as a case study we focus on Guo & Yannelis (2022), where an analysis of belief-free full implementation in the spirit of Bergemann & Morris (2009) and Bergemann & Morris (2011) is performed in presence of possible coalitions, with the understanding that a coalition can be an example of a conditioning event, and—incidentally—it is also an example of how our topology-free construction with unrestricted conditioning events can be a foundation for ‘robustness’ endeavours in the spirit of Bergemann & Morris (2005) when in presence of conditioning events and—in particular—counterfactual reasoning.

To see this, the following is—modulo notation and terminology—the framework of Guo & Yannelis (2022, Section 2, pp.555–556), where a *payoff environment* is a tuple

$$\mathcal{E} := \langle I, X, A, (\Theta_i, u_i)_{i \in I} \rangle$$

with I being a set of *agents*, X a set of *feasible deterministic outcomes*, $A := \Delta(X)$ a set of *feasible outcomes*, whereas Θ_i is the space of payoff types of player i , with $\Theta := \prod_{j \in I} \Theta_j$, and $u_i : A \times \Theta \rightarrow \mathfrak{R}$ is her Bernoulli (unique up to positive affine transformation) utility function. Given a payoff environment \mathcal{E} , an information structure appended on \mathcal{E} is a tuple $\mathcal{Y} := \langle (Y_i, \varsigma_i, \varpi_i)_{i \in I} \rangle$, where, for every $i \in I$, Y_i is agent i ’s *information space*, $\varsigma_i : Y_i \rightarrow \Theta_i$ is her *payoff type function*, and $\varpi_i : Y_i \rightarrow \Delta(Y_{-i})$ is her *belief function*.⁵⁵ Finally, to capture the presence of coalitions and corresponding beliefs of a player i , given that $\mathcal{I} := 2^I \setminus \{\emptyset\}$ denotes the set of *coalitions* and that

$$\varpi_{i,J}(y_i)(y_{S \setminus \{i\}}) := \text{marg}_{Y_{S \setminus \{i\}}} \varpi_i(y_i)(y_{-i}),$$

where $J \in \mathcal{I}$ with $i \in J$ and $y := (y_i)_{i \in I} \in Y$ (with marg denoting the marginal operator as canonically defined), a *belief revising rule* specifies a posterior belief $\varpi_i(y_i)(y_{-i}) \in \Delta(Y_{-i})$ with $\varpi_{i,J}(y_i)(y_{J \setminus \{i\}}) = 1$ whenever $\varpi_{i,J}(y_i)(y_{J \setminus \{i\}}) = 0$, where the posterior belief is defined via the Bayes rule (i.e., Axiom (C3) in Definition 2.1) whenever $\varpi_{i,J}(y_i)(y_{J \setminus \{i\}}) > 0$.

Now, it turns out that our construction in Section 4.2 allows to obtain the topology free universal type structure in presence of unrestricted conditioning events $\mathcal{T}^* := \langle (T_i^*, \beta_i^*)_{i \in I} \rangle$ for the setting just described, where an arbitrary type structure \mathcal{T} is essentially equivalent to an information

⁵⁴Alternatively called “Extensive-Form Rationalizability” (as, for example, in Battigalli & Siniscalchi (2002, Definition 5, p.373)). Here, we adopt a terminology used for example in Battigalli & Siniscalchi (2003), which has the virtue of distinguishing this one from other forms of Rationalizability that can be implemented in the analysis of dynamic games represented in their extensive-form (i.e., Initial (or Weak) Rationalizability *à la* Ben-Porath (1997), Backward Rationalizability *à la* Penta (2015), or Forward and Backward Rationalizability *à la* Meier & Perea (2023)).

⁵⁵In Guo & Yannelis (2022, Section 2, p.556), the authors actually use the expression “type structure” to refer to this object and use T_i to denote the type space of an arbitrary player i . Here, we opt to call this object “information structure” with Y_i being agent i ’s *information space* to avoid an overloading of the previous expressions and notation.

structure \mathcal{Y} given appropriate restrictions. Thus, in particular, given that $\mathcal{C}_I := \mathcal{I}$ with $\mathbf{P}_I := \mathbf{K}_I \times \mathbf{K}_\Theta \times \mathbf{Proj}_{-i}$, we need to define the new functor $\mathbf{C} := \Delta^{\mathcal{C}_I} \mathbf{P}_I$, i.e.,

$$\mathbf{C} := \Delta^{\mathcal{C}_I} \mathbf{P}_I = \Delta^{\mathcal{C}_I} (\mathbf{K}_I \times \mathbf{K}_\Theta \times \mathbf{Proj}_{-i}),$$

where the appropriate definitions regarding the way in which the functor acts on objects and morphisms follow the path described in [Section 4.2](#), which, as a result, allows us to obtain a belief function

$$\beta_i^* := (\beta_{i,J}^*)_{J \in \mathcal{I}} : T_i \rightarrow \Delta^{\mathcal{I}}(I \times \Theta \times T_{-i}).$$

Given the steps sketched above, it remains to carve out (in the spirit of [Section 5.1](#)) the resulting structure, a belief-closed type structure with the desired properties which is universal in its own rights given these properties, where—in particular—we have to impose that β_i^* is such that $\text{marg}_J \beta_{i,J}^*(t_i) = \delta_J$ and $\text{marg}_{\Theta_i} \beta_{i,J}^*(t_i) = \delta_{\{\theta_i\}}$ for a $\theta_i \in \Theta_i$, for every $t_i \in T_i$ and $J \in \mathcal{I}$, where δ_J and $\delta_{\{\theta_i\}}$ denote the Dirac measure as canonically defined. Indeed, this delivers exactly what we want in light of the obvious relation between $\mathcal{Y} := \langle (Y_i, \varsigma_i, \varpi_i)_{i \in I} \rangle$ and an arbitrary belief-closed type structure $\mathcal{T} := \langle (T_i, \beta_i)_{i \in I} \rangle$ with β_i defined along the lines of β_i^* above.

6. DISCUSSION

6.1 Relation to the Literature

6.1.1 Relation to [Heifetz & Samet \(1998\)](#)

Concerning the construction in [Heifetz & Samet \(1998, Section 5\)](#), there is a tight relation between the construction performed here and the one delivered therein. Indeed, the importance of the construction in [Heifetz & Samet \(1998, Section 5\)](#) cannot be emphasized enough, since, starting from the usage of a singleton set, it is essentially coalgebraic in nature. Indeed, it is possible to find a tight link between all the results obtained [Heifetz & Samet \(1998, Section 5\)](#) and those in [Viglizzo \(2005b\)](#). However, as it has already been pointed out in [Section 1.1](#), it is important to stress that the construction in [Heifetz & Samet \(1998, Section 5\)](#) does *not* deliver all the universality properties at once, which is the reason why we employ coalgebraic tools in the present work.⁵⁶

6.1.2 Relation to [Viglizzo \(2005b\)](#)

Regarding [Viglizzo \(2005b\)](#), it is important to emphasize two—somewhat related—points. The first one is that the result (and corresponding proof) in [Viglizzo \(2005b\)](#) is actually more general than [Proposition 2](#). Indeed, [Viglizzo \(2005b, Theorem 1, p.404\)](#) states that there exists a terminal Φ -coalgebra, for *every* endofunctor Φ on \mathbf{Meas} which is build from the identity, the constant, and an appropriately defined Δ functor (as $\Delta(X) := \Delta(X)$, for every $X \in \text{Ob}(\mathbf{Meas})$), and Δf as the image measure of a morphism f belonging to \mathbf{Meas} with a closure over binary products and coproducts, i.e., disjoint unions. Thus, for example, this result tells us that there exists a terminal Φ -coalgebra for the—admittedly cumbersome—functor $\Phi := \mathbf{Id} \times \mathbf{K}_M \times \Delta(\mathbf{Id} \sqcup \mathbf{K}_M)$. Now, our [Proposition 2](#) does not deliver anything like that, but it is rather tailor-made to deal with type structures with conditioning events, which considerably changes the nature of the problem, the point being that, for example, a minimal departure from the theoretical problem we faced such as the expression $\Delta^{\mathcal{C}_M}(\mathbf{Id})$ would actually be ill-defined. Nonetheless, in the spirit of the minimal adaptation made in [Section 5.3](#) of the path sketched in [Section 4.2](#), we can obtain a result concerning the flexibility of the form of the functor we want to focus on as in [Viglizzo \(2005b\)](#), leveraging on the definition of ζ as $\zeta := \vec{\zeta} \circ \zeta_+$ and always taking into account the peculiarities of the functor $\Delta^{\mathcal{C}_M}$ introduced here. In particular, this is a byproduct of the fact that the path we follow and the proofs of the results we deliver in [Section 4.1](#) are (modulo notation) those in [Viglizzo \(2005b\)](#),⁵⁷ with the notable exception of [Lemma C.1](#), which—even if along the lines of the corresponding [Viglizzo \(2005b, Lemma 8, pp.402–403\)](#)—deals with the presence of conditioning events and, as a result, shows how the crucial new technical element of the present work lies in identifying the peculiarities that arise in presence of conditioning events, and of [Lemma 5](#), which ‘fixes’—contrary, as pointed out above, to [Viglizzo \(2005b\)](#)—the ‘form’ of the functor we are interested in.

⁵⁶See also [Section 6.2.3](#) with respect to this point.

⁵⁷That we rewrite here with our notation for self-containment purposes.

Now, the generality of [Viglizzo \(2005b\)](#) is not a peculiarity of this very paper, but rather of the nature of the problems addressed in the stream of literature to which the paper belongs, which can be captured by the question if it is possible to build a terminal coalgebra for a given collection of endofunctors built from ‘more basic’ endofunctors on a given category. Thus, this is related to the second point, namely, that this paper and the literature to which it belongs tell us something about results from interactive epistemology at large. As an example, we know from [Brandenburger \(2003, Proposition 1, p.32\)](#) that there exists no belief-complete interactive—possibility—structure of the form $\mathcal{P} := \langle (T_i, \wp_i)_{i \in I} \rangle$, where $\wp_i : T_i \rightarrow \mathcal{P}(\Theta \times T_{-i})$. However, strictly speaking, this is *neither* a consequence of the functors upon which the functor \mathcal{P} acts (in the spirit of [Section 4.1](#)), e.g., \mathcal{P} -coalgebras for a functor of the form $\mathcal{P}(\mathbf{K}_\Theta \times \mathbf{Id})$, *neither* it is a result specific of interactive structures built on the functor \mathcal{P} (in the spirit of [Section 4.2](#)), i.e., \mathcal{P} -coalgebras for a functor of the form $\mathcal{P}(\mathbf{K}_\Theta \times \mathbf{Proj}_{-i})$, but rather it is a more basic issue concerning the very functor \mathcal{P} , as pointed out in [Section 2.2](#), which shows us that the impossibility runs deeper than the product structure we typically deal with in interactive epistemology, i.e., the problem is *functorial*.

6.1.3 Relation to [Heinsalu \(2014\)](#)

[Heinsalu \(2014\)](#) is an important paper in the literature on the construction of large type structures: beyond addressing the conceptually challenging problem of such a construction for the case of unawareness in a static setting, this is the first paper belonging to the game-theoretical literature where coalgebraic methods are employed. Thus, it is essential to identify how this paper is different from the present one. Now, bypassing the fact that [Heinsalu \(2014\)](#) deals with a static setting in presence of unawareness, whereas our focus is on a setting with conditioning events without unawareness, the major difference between these two papers lies in the background work. [Heinsalu \(2014\)](#) is crucially based on the background work and the results obtained in [Viglizzo \(2005b\)](#) and the relevant functor for the enterprise is ready-made. On the contrary, here, we have to identify the properties of the spaces we are dealing with along with the appropriate functors to address the problem. Indeed, a major point of the present work lies in identifying the peculiarities of product conditional measurable spaces (in general) and product conditional measurable spaces sharing a family of conditioning events (in particular), which leads to the definition of a new—tailor-made for the problem at hand—functor, i.e., the functor $\Delta^{\mathbf{C}_M}$ upon which the functor $\bar{\mathbf{T}} := \Delta^{\mathbf{C}_M} \mathbf{P}_M$ is based. As such, the present work, by additionally spelling out the details of the—relevant—proofs from [Viglizzo \(2005b\)](#) can prove to be a useful introduction to categorical and coalgebraic methods.

6.2 Relation to Universality Notions

6.2.1 On the Notion of Kolmogorov’s Consistency

It is rather natural to ask ourselves why all of this was needed after all. Thus, in order to answer this question, we need to frame [Definition 4.6](#) in a more general way. In particular, given an index set Λ treated as a category⁵⁸ and an arbitrary category \mathbf{C} , a *diagram* \mathbf{D} is a functor from Λ to \mathbf{C} . Hence, a cone⁵⁹ on \mathbf{D} is an object $A \in \text{Ob}(\mathbf{C})$ together with a family of maps $(A \xrightarrow{f_\lambda} \mathbf{D}_\lambda)_{\lambda \in \Lambda}$, such that the following diagram commutes

$$\begin{array}{ccc} & A & \\ f_\lambda \swarrow & & \searrow f_{\lambda'} \\ \mathbf{D}_\lambda & \xleftarrow{\mathbf{D}_u} & \mathbf{D}_{\lambda'} \end{array}$$

for every $\lambda' \xrightarrow{u} \lambda$. Now, a *limit* L is a cone $(L \xrightarrow{f_\lambda} \mathbf{D}_\lambda)_{\lambda \in \Lambda}$ with the property that there exists a unique morphism $\bar{f} : A \rightarrow L$ such that $\pi_\lambda \circ \bar{f} = f_\lambda$, for every $\lambda \in \Lambda$ (with π_λ denoting—*as usual*—the projection operator), where it is immediate to observe that this is exactly what we

⁵⁸Technically, this is a *small category*, i.e., a category whose both objects and morphisms form a set, i.e., not a class (see [Leinster \(2014, Chapter 3.2, p.75\)](#) for the notion of small category and [Jech \(2006, Chapter 1, pp.5–6\)](#) for the notion of class in set theory). Thus, the set \mathbb{N} can be taken as the index set.

⁵⁹See [Leinster \(2014, Chapter 5.1, p.118\)](#) for the definition of cone.

introduced in Section 4.1.1 framed in more general terms. Finally, a functor Φ preserves limits if, given a limit $(L \xrightarrow{f_\lambda} \mathbf{D}_\lambda)_{\lambda \in \Lambda}$, we have that $(\Phi L \xrightarrow{\Phi f_\lambda} \Phi \mathbf{D}_\lambda)_{\lambda \in \Lambda}$ is a limit.

The problem lies in the fact that the functor $\Delta^{\mathcal{C}_M}$ does not preserve limits in the form of ω^{op} -chains, i.e., by employing a common terminology in the literature, it is not ω^{op} -continuous. Indeed, this is an immediate consequence of Viglizzo (2005b, Section 2.2, p.399) in light of the fact that product conditional measurable spaces are a ‘special’ case of measurable spaces, i.e., and $\Delta(M) = \Delta^{\mathcal{C}_M}(M)$ for an arbitrary M when \mathcal{C}_M is a singleton. Of course, the point made in Viglizzo (2005b, Section 2.2, p.399) is tightly related to the impossibility of obtaining an analog of the Kolmogorov’s Extension Theorem⁶⁰ for an infinite product of measurable spaces, as shown by Sparre Andersen & Jessen (1948), Dieudonné (1948, Section 7, p.42), and Halmos (1950, Chapter IX.49(3), p.214), which is—in turn—related to the result obtained in Heifetz & Samet (1999) concerning the fact that there are coherent infinite hierarchies of beliefs that are not types, i.e., when we do not start from topological assumptions, it is possible to obtain a coherent infinite hierarchy of beliefs such that there exists no extension over the infinite product (which would be essentially the type, as employed in economics). In other words, by letting C denote the space of coherent infinite hierarchies of beliefs over a common domain of uncertainty Θ with T^* the corresponding space of *all* types, we have $T^* \subseteq C$, with strict inclusion $T^* \subset C$ when we work without topological assumptions.⁶¹ Relatedly, it is important to point out that it has been shown in Schubert (2009) that starting with appropriate topological assumptions (and relying on a suitable version of the Kolmogorov’s Extension Theorem), it is possible to build an endofunctor that preserves limits in the form of ω^{op} -chains and allows to obtain the corresponding terminal coalgebra via the canonical hierarchical construction presented in Section 4.1.1.⁶²

6.2.2 On Belief-Completeness & Relation to Meier (2012)

For the ‘standard’ case without conditioning events, the isomorphic nature of the belief functions in the construction in Heifetz & Samet (1998) along with—as a corollary—its belief-completeness has been established for the first time in Meier (2012, Theorem 4, p.29),⁶³ where the author builds an infinitary probability logic to capture reasoning in type structures.⁶⁴ Thus, to relate this work to the present one, it is worthwhile to recall the path taken in Meier (2012) to obtain the result. Hence, Meier (2012) starts by addressing belief structures, where, given a measurable space (Θ, Σ_Θ) , a *belief structure*⁶⁵ (as in Meier (2012, Definition 7, p.11), in its full generality) is a tuple $\mathcal{B} := \langle \Omega, \theta, (\tau_i)_{i \in I} \rangle$ where $\theta \in \Theta^\Omega$ is a measurable function and $\tau_i \in [\Delta(\Omega)]^\Omega$ is measurable, for every $i \in I$. Thus, first of all, completeness (in the logical sense as in Blackburn et al. (2001, Chapter 4)) is established for the logical system introduced to deal with belief structures; then, it is established the terminality of the constructed structure canonically built from appropriate formulas; finally, it is showed that the canonical structure has a product structure (in Meier (2012, Proposition 4(3), p.28)) and it is belief-complete (in Meier (2012, Theorem 4, p.29)).

In light of the previous paragraph, three points are in order. The first one is that, in this work, we obtain the belief-completeness of the universal type structure \mathcal{T}^* for free, as a corollary of Lambek’s lemma. As a matter of fact, we see this as a positive side of performing the endeavour in a coalgebraic framework, which can be exploited by practitioners working on the existence of large interactive structures in topology-free settings to obtain all relevant results at once. The second one is that, whereas we built the terminal \mathbf{T} -coalgebra via so-called final sequences, it is possible to obtain the same result via tools from coalgebraic modal logic, as done in Moss & Viglizzo

⁶⁰See Aliprantis & Border (2006, Theorem 15.26, p.522).

⁶¹In particular, see Heifetz & Samet (1999, Section 5).

⁶²In particular, Schubert (2009) works in the subcategory **SB** of **Meas** comprised of standard Borel spaces (i.e., those measurable spaces (X, \mathcal{B}_X) such that there exists a Polish topology τ_X on X with \mathcal{B}_X being the family of Borel sets generated by τ_X) with the subprobability endofunctor S , where a subprobability measure μ on a standard Borel space (X, \mathcal{B}_X) is a σ -additive functional on \mathcal{B}_X such that $\mu(X) \in [0, 1]$ (see also Moss (2011, Section 2.4, p.33)).

⁶³The first version of this (in the words of Aumann & Heifetz (2002, Footnote 14, p.1674), where there is also a brief—nonetheless informative—description of part of the results therein) “*beautiful, path-breaking paper*” goes back to 2001, thus establishing the result before Moss & Viglizzo (2004).

⁶⁴See also Heifetz (1997), Aumann (1999b), and Heifetz & Mongin (2001).

⁶⁵See Mertens & Zamir (1985, Definition 2.2, p.4) or Maschler et al. (2013, Chapter 10).

(2004),⁶⁶ where it would be interesting to see the relation between these logics and those as the one obtained in Meier (2012). Finally, obtaining the belief-completeness of our structure is one of the differences between the present work and Fukuda (2021, Section C.1, Online Appendix), where only terminality is established, the other being that the focus in that work is on belief structures.⁶⁷

6.2.3 On the Notion of Universality

First of all, we want to emphasize one point about the terminology we use, namely our—somewhat non-standard—definition of universality with respect to how the term is used in the literature on large interactive structures. As mentioned in Section 3.4, we extend a terminology that goes back to Siniscalchi (2008, p.93) according to which a type structure is universal if terminal and belief-complete. However, starting from Mertens & Zamir (1985, Theorem 2.9, pp.7–8),⁶⁸ the notion of universal type structure has been associated with the idea that any other type structure can be uniquely embedded into the universal one. In other words, the term “universal” has been used as a synonym for “terminal”. Interestingly, and somewhat reconciling the usage of both terms for the same object, it should be pointed out how this very definition of universality fits the categorical usage of the word “universal” in the opening quotation of this work by Dieudonné (1989): every other type structure can indeed be built from the *universal* one.

In second place, it is important to address the relation between the notion of terminality and its connection to the notion of universality—as described above—in its categorical interpretation and the other notions of belief-completeness and non-redundancy. Thus, in particular, it is natural to ask ourselves if terminality actually *implies* all these properties.⁶⁹ The answer is: *it depends!* Indeed, the problem lies in how we look at this issue. If we approach it from a purely mathematical standpoint, terminality does not necessarily implies all the other properties due to what happens in the topology-free case, where the construction of the terminal type structure as performed in Heifetz & Samet (1998) does *not* deliver automatically belief-completeness (obtained—as pointed out in Section 1.1 and Section 6.2.2—for the first time in Meier (2012)). However, it is also important to recognize that, exactly due to ‘categorical’ reasons, terminality from a conceptual standpoint does ‘imply’ all the other properties and, once certain tools are employed (for example, in the topology-free case without conditioning events, infinitary probability logic as in Meier (2012) or coalgebraic methods as in Viglizzo (2005b)), indeed it turns out that the terminal type structure *is* the universal one. In particular, with respect to this point, it could be possible to introduce a new notion of universality, call it “*Categorical Universality*”, which would correspond to the idea of a type structure being terminal (and unique up to type isomorphism), non-redundant, and such that all the belief functions are *isomorphisms* in the relevant category. Indeed, this would capture exactly the properties we obtain via the coalgebraic canonical construction, with belief-completeness being an immediate consequence.

Now, it seems that the issues pointed out above strongly support the usage of coalgebraic methods for the construction of large interactive structures in light of the fact that, via these tools, terminality *implies* all the properties exactly as in the case of the topological constructions, where the usual canonical construction⁷⁰ of a type structure that is terminal automatically delivers its universality. In particular, in the spirit of what pointed out in Section 6.2.1, it is worth emphasizing that the coalgebraic canonical construction presented in Section 4.1.1 can be—in principle—delivered in both the topological and the topology-free case,⁷¹ thus, turning out to be a bridge between constructs obtained via procedures at the present stage seemingly different.⁷²

⁶⁶See also Moss & Viglizzo (2006). Regarding coalgebraic logic, see Jacobs (2001), Kurz (2001), and Rößiger (2001) or Jacobs (2017, Chapter 6.5) for a textbook presentation.

⁶⁷See also Di Tillio et al. (2014).

⁶⁸For example, see also Heifetz & Samet (1998, Definition 3.3, p.331).

⁶⁹We are extremely grateful to an anonymous referee for having raised this point.

⁷⁰With the caveat that, in the topological case, there are essentially two canonical constructions as in the terminology introduced in Battigalli et al. (Work in Progress): i.e., the *bottom-up constructions*, with the coherency requirement imposed at the outset in the spirit of Mertens & Zamir (1985), and the *top-down constructions*, where first infinite hierarchies of beliefs are constructed and then the coherency requirement is imposed on them, like in Brandenburger & Dekel (1993).

⁷¹See Heinsalu (2014, Section 2, p.257) for a similar point.

⁷²And it is worth mentioning with respect to the notion of Categorical Universality set forth above that homeomorphisms are—indeed—*isomorphisms* in the category **Top** of topological spaces and continuous functions between them.

6.3 On Technical Aspects of the Construction

6.3.1 On Type Structures

Regarding our definition of type structure as in [Definition 3.1](#), it should be observed that we could have defined the belief function of agent i as $\beta_i : T_i \rightarrow \Delta^{\mathcal{C}_\Theta}(\Theta \times T)$, for every $i \in I$, as in [Heifetz & Samet \(1998, Definition 3.1, pp.329–330\)](#). With such a definition, it is typically imposed the additional requirement that $\text{marg}_{T_i} \beta_{i,C}(t_i) = \delta_{t_i}$, for every $t_i \in T_j$ and $C \in \mathcal{C}_\Theta$,⁷³ with marg denoting the marginal operator as canonically defined and δ_{t_i} denoting the Dirac measure on t_i . It has to be observed that this property is not necessary for the construction: [Heifetz & Mongin \(2001, Section 5, p.41\)](#) and [Meier \(2012, Definition 8, p.12\)](#) distinguish type structures which possess this property from those which do not,⁷⁴ where in their terminology a type structure that satisfies this requirement is called a “*Harsanyi type structure*”.⁷⁵

6.3.2 On Type Morphisms

Concerning our definition of type morphism, as it is widespread in the literature, given an arbitrary conditional measurable space $(\Theta, \Sigma_\Theta, \mathcal{C}_\Theta)$ and a type structure \mathcal{T} , we defined it in [Definition 3.2](#) by introducing a function $\vartheta_0 := \mathbf{id}_\Theta$. Interestingly, this is not *explicitly* present in the notion of \mathbf{T} -coalgebra morphism. Now, a similar issue is present in [Moss & Viglizzo \(2004\)](#), as emphasized in [Moss & Viglizzo \(2004, Section 2, pp.284–285\)](#), where the authors point out that their notation remains silent regarding the underlying (common) domain of uncertainty. Of course, our notation does not—and cannot—remain silent, since emphasizing the common domain of uncertainty is crucial to properly deal with product conditional measurable spaces sharing a given set of conditioning events. Thus, as a matter of fact, our proof takes care of the existence of the function ϑ_0 in light of the definition of \mathbf{P}_Θ , which is based on a binary product involving \mathbf{K}_Θ , that—in turn—induces a morphism \mathbf{id}_Θ .

6.3.3 On the Family of Conditioning Events

The present construction does not put any restriction on the conditioning events belonging to a given conditional measurable space $(M, \Sigma_M, \mathcal{C}_M)$. Thus, it is important to make a comparison with the original construction obtained in [Battigalli & Siniscalchi \(1999, Section 2\)](#) by paying special attention to the assumptions made there and the rationale behind them. In particular, in [Battigalli & Siniscalchi \(1999, Section 2.1, p.191\)](#), it is assumed that, given a conditional measurable space $(M, \Sigma_M, \mathcal{C}_M)$, M is a Polish space, Σ_M is its Borel σ -algebra, and the events in \mathcal{C}_M are clopen (i.e., closed and open) and at most countable.

Regarding the clopeness of the conditioning events in \mathcal{C}_M , this assumption is used in [Battigalli & Siniscalchi \(1999, Proof of Lemma 1, p.224\)](#) to establish that $\Delta^{\mathcal{C}_M}(M)$ is a closed subset of $[\Delta(M)]^{\mathcal{C}_M}$. Concerning the cardinality assumption, assuming an at most countable family of conditioning events is crucial in [Battigalli & Siniscalchi \(1999, Proof of Proposition 1, pp.196–197\)](#), where it is used to prove the existence of—using our notation—a homeomorphism $f : H_c \rightarrow \Delta^{\mathcal{C}_M}(M \times H)$, where H is the set of infinite hierarchies of beliefs not necessarily coherent, whereas H_c is the set of infinite hierarchies of beliefs that are coherent. In particular, this assumption is needed to establish, via arguments from [Kechris \(1995, Section 11.A, p.68\)](#) concerning Borel sets, that every n^{th} -order hierarchy of CPSs belonging to a CPS over $\Delta^{\mathcal{C}_M}(M \times H)$ satisfies Axiom (C3) from [Definition 2.1](#).

It is immediate to observe that, while the clopeness assumption simply has no bite in our topology-free framework, the cardinality assumption is not needed to establish our result. Hence, our construction does not put any restriction on the nature of the conditioning events.⁷⁶

⁷³In particular, this is actually Condition (3) in [Heifetz & Samet \(1998, Definition 3.1, pp.329–330\)](#).

⁷⁴See also the discussion of this point in a coalgebraic context in [Viglizzo \(2005a, Chapter 7.1, pp.79–81\)](#).

⁷⁵With the caveat that these authors employ the word “space” instead of “structure”.

⁷⁶We are extremely grateful to an anonymous referee for having questioned the need of the countability assumption we had in a previous version of the paper. In doing so, we realized we could go back to work with ‘unrestricted’ conditioning events in the spirit of [Guarino \(2017\)](#) and [Fukuda \(2021\)](#) (and, with respect to this point, it is worthwhile mentioning that neither cardinality nor topological assumptions are made in the original [Rényi \(1955\)](#)).

APPENDICES

A. CATEGORY THEORY BACKGROUND

In this section, for self-containment reasons, we provide an introduction to those elements of category theory that we need for our purposes. Thus, first of all, we define what a category actually is.

Definition A.1 (Category). *A category \mathbf{C} consists of the following data:*

- a collection of objects, denoted by $\text{Ob}(\mathbf{C})$;
- for every $A, A' \in \text{Ob}(\mathbf{C})$, a collection of morphisms (or arrows) from A to A' , denoted by $\mathbf{C}(A, A')$;
- for every $A, A', A'' \in \text{Ob}(\mathbf{C})$, a composition law

$$\mathbf{C}(A, A') \times \mathbf{C}(A', A'') \rightarrow \mathbf{C}(A, A''),$$

where the composite of $f \in \mathbf{C}(A, A')$ and $f' \in \mathbf{C}(A', A'')$ is denoted $f' \circ f$;

- for every $A \in \text{Ob}(\mathbf{C})$, a morphism $\text{id}_A \in \mathbf{C}(A, A)$, called the identity on A .

These data satisfy the following axioms:

- associativity: for every $f \in \mathbf{C}(A, A')$, $f' \in \mathbf{C}(A', A'')$, $f'' \in \mathbf{C}(A'', A''')$,

$$(f'' \circ f') \circ f = f'' \circ (f' \circ f);$$

- identity: for every $f \in \mathbf{C}(A, A')$,

$$f \circ \text{id}_A = f = \text{id}_{A'} \circ f.$$

Having introduced the notion of category, it is typical to provide actual examples of categories, since it could be stated that one of the major points of embracing a ‘categorical’ approach is the fact that it is possible to identify properties shared by seemingly exceedingly different constructs.

Example A.1 (Examples of Categories). We now present two examples of categories that show how flexible [Definition A.1](#) actually is.

- The category **Set** is the category whose objects are sets and whose morphisms are functions between sets. Relevant for our purposes, given $A, A' \in \text{Ob}(\mathbf{Set})$, we have $A \times A' \in \text{Ob}(\mathbf{Set})$.
- The category **Top** is the category whose objects are topological spaces and whose morphisms are continuous functions between topological spaces. ◇

Definition A.2 (Subcategory). *Given a category \mathbf{C} , a subcategory \mathbf{S} of \mathbf{C} consists of:*

- a subclass $\text{Ob}(\mathbf{S})$ of $\text{Ob}(\mathbf{C})$ and
- a subclass $\mathbf{S}(A, A')$ of $\mathbf{C}(A, A')$

such that \mathbf{S} is closed under composition and identities. A subcategory is full if $\mathbf{S}(A, A') = \mathbf{C}(A, A')$, for every $A, A' \in \text{Ob}(\mathbf{S})$.

Given two categories, it is possible to have morphisms between the two. The next definition captures exactly what are the properties that these morphisms, called functors, need to satisfy in order to be ‘well-behaved’.

Definition A.3 (Functor). *Given two categories \mathbf{C} and \mathbf{C}' , a functor $\Phi : \mathbf{C} \rightarrow \mathbf{C}'$ between \mathbf{C} and \mathbf{C}' consists of:*

- a morphism $\text{Ob}(\mathbf{C}) \rightarrow \text{Ob}(\mathbf{C}')$, written $A \mapsto \Phi(A)$;
- for every $A, A' \in \text{Ob}(\mathbf{C})$, a morphism

$$\mathbf{C}(A, A') \rightarrow \mathbf{C}'(\Phi(A), \Phi(A')),$$

written $f \mapsto \Phi(f)$.

These data satisfy the following axioms:

- AC) $\Phi(f' \circ f) = \Phi(f') \circ \Phi(f)$, with $f \in \mathbf{C}(A, A')$ and $f' \in \mathbf{C}(A', A'')$,
- AI) $\Phi(\text{id}_A) = \text{id}_{\Phi(A)}$, for every $A \in \text{Ob}(\mathbf{C})$.

The notion of functor captures the crucial idea of *functoriality*. Thus, given two categories \mathbf{C} and \mathbf{C}' , if we define a morphism Φ between them and we prove that Φ actually is a functor, i.e., it satisfies Axioms (AC) and (AI), then it turns out that Φ does not simply act on the elements $\text{Ob}(\mathbf{C})$, but also on its morphisms in an appropriate way.⁷⁷

Example A.2 (Product Categories & Projection Functors). Given a *product* category $\prod_{\lambda \in \Lambda} \mathbf{C}_\lambda$, the *projection functor* \mathbf{Proj}_λ is the functor defined as

- $(A_\lambda)_{\lambda \in \Lambda} \mapsto A_\lambda$, for every $(A_\lambda)_{\lambda \in \Lambda} \in \text{Ob}(\prod_{\lambda \in \Lambda} \mathbf{C}_\lambda)$,
- $(f_\lambda)_{\lambda \in \Lambda} \mapsto f_\lambda$, for every $(f_\lambda)_{\lambda \in \Lambda} \in \mathbf{C}_\lambda((A_\lambda)_{\lambda \in \Lambda}, (A'_\lambda)_{\lambda \in \Lambda})$,

for every $\lambda \in \Lambda$. Now, let $\mathbf{Set}^2 := \mathbf{Set} \times \mathbf{Set}$ and assume that A, A' are exactly the same sets considered in [Example A.1](#). It is important to observe that $A \times A' \in \text{Ob}(\mathbf{Set})$ is not the same as $A \times A' \in \text{Ob}(\mathbf{Set}) \times \text{Ob}(\mathbf{Set})$ since the two expressions capture objects belonging to different categories. \diamond

In this paper, we focus on a specific class of functors, namely, those functors that map a category to itself.

Definition A.4 (Endofunctor). Given a category \mathbf{C} , an endofunctor on \mathbf{C} is a functor $\Phi : \mathbf{C} \rightarrow \mathbf{C}$.

Example A.3 (Three Endofunctors). We now provide three examples of endofunctors: the first two are phrased in rather abstract terms and are extensively used in our construction, while the third is an example of a functor in the category \mathbf{Set} .

- Given an arbitrary category \mathbf{C} , the *identity (endo)functor* $\text{Id} : \mathbf{C} \rightarrow \mathbf{C}$ is defined as $\text{Id}(A) := A$, for every $A \in \text{Ob}(\mathbf{C})$, and $\text{Id}(f) := f$, for every $f \in \mathbf{C}(A, A')$, with $A, A' \in \text{Ob}(\mathbf{C})$;
- Given an arbitrary category \mathbf{C} , the *constant endofunctor given X*, denoted $\mathbf{K}_X : \mathbf{C} \rightarrow \mathbf{C}$, is defined as $\mathbf{K}_X(A) := X$, for every $A \in \text{Ob}(\mathbf{C})$, and $\mathbf{K}_X(f) := \text{id}_X$, for every $f \in \mathbf{C}(A, A')$, with $A, A' \in \text{Ob}(\mathbf{C})$.
- Building on the category \mathbf{Set} , we now provide an example of an endofunctor on \mathbf{Set} . Thus, let $\mathcal{P} : \mathbf{Set} \rightarrow \mathbf{Set}$ be defined as:

- regarding objects, $\mathcal{P}(A) := \mathcal{P}(A)$, for every $A \in \text{Ob}(\mathbf{Set})$, with \mathcal{P} denoting the power set;
- regarding morphisms,

$$\mathcal{P}(f) : \mathcal{P}(A) \rightarrow \mathcal{P}(A')$$

such that $\mathcal{P}(f)(B) := f(B)$, for every $f \in \mathbf{Set}(A, A')$ and $B \in \mathcal{P}(A)$.

The resulting morphism is the so-called *power set (endo)functor*.⁷⁸ \diamond

⁷⁷Regarding this notion, see the discussion (and the examples) in [Jacobs & Rutten \(1997, Section 4\)](#).

⁷⁸For completeness, this is called the *covariant* power set functor: see [Leinster \(2014, Chapter 1.2, p.22\)](#) for an explanation of this terminology.

Having established the setting above, it is possible to define when two objects in a given category are—in a sense to be precisely defined—the same. The two definitions that follow accomplish exactly this objective.

Definition A.5 (Isomorphism). *Given a category \mathbf{C} and objects $A, A' \in \text{Ob}(\mathbf{C})$, $f \in \mathbf{C}(A, A')$ is an isomorphism if there exists a morphism $f' \in \mathbf{C}(A', A)$ such that the following diagram*

$$\begin{array}{ccccc} & & & \xrightarrow{f \circ f' = \text{id}_{A'}} & \\ & & & \nearrow & \\ A & \xrightarrow{f} & A' & \xrightarrow{f'} & A & \xrightarrow{f} & A' \\ & & & \searrow & \\ & & & \xrightarrow{f' \circ f = \text{id}_A} & \end{array}$$

commutes.

Definition A.6 (Isomorphic Objects). *Given a category \mathbf{C} and objects $A, A' \in \text{Ob}(\mathbf{C})$, A and A' are isomorphic, denoted $A \cong A'$, if there exists an isomorphism between A and A' .*

Finally, it is the existence of a special kind of objects in a category, namely, the so-called terminal ones, that plays a major role in this paper.

Definition A.7 (Terminal Object). *Given a category \mathbf{C} , an object $\mathbf{1} \in \text{Ob}(\mathbf{C})$ is terminal if for every $A \in \text{Ob}(\mathbf{C})$ there exists a unique morphism $! \in \mathbf{C}(A, \mathbf{1})$ such that $! : A \rightarrow \mathbf{1}$, i.e., $A \xrightarrow{!} \mathbf{1}$, for every $A \in \text{Ob}(\mathbf{C})$.*

B. MEASURE-THEORETIC RESULTS

We recall that, given a measurable space (M, Σ_M) , a π -system is a nonempty family of subsets $\mathcal{D} \subseteq \mathcal{P}(M)$ that is closed under finite intersections. The next result is an immediate consequence of Dynkin's π - λ Lemma⁷⁹ applied to all the conditioning events in \mathcal{C}_M .

Lemma B.1. *Given a product conditional measurable space $(M \times X, \Sigma_M \otimes \Sigma_X, \mathcal{C}_M)$, where $\Sigma_M \otimes \Sigma_X := \sigma(\mathcal{D})$ with \mathcal{D} an arbitrary π -system, and two CPSs $\nu, \nu' \in \Delta^{\mathcal{C}_M}(M \times X)$, if $\nu(D|C) = \nu'(D|C)$, for every $D \in \mathcal{D}$ and $C \in \mathcal{C}_M$, then $\nu(E|C) = \nu'(E|C)$, for every $E \in \sigma(\mathcal{D})$ and $C \in \mathcal{C}_M$.*

The next remark is an analog of [Viglizzo \(2005b, Lemma 2, p.396\)](#), which in turn is in the spirit of [Heifetz & Samet \(1998, Equation 3.2, p.330\)](#).

Remark B.1. *Given a conditional measurable space $(M, \Sigma_M, \mathcal{C}_M)$, two product conditional measurable spaces $(M \times X, \Sigma_M \otimes \Sigma_X, \mathcal{C}_M)$ and $(M \times Y, \Sigma_M \otimes \Sigma_Y, \mathcal{C}_M)$ sharing \mathcal{C}_M , and a measurable function $f \in (M \times Y)^{(M \times X)}$,*

$$\gamma_C^p(f^{-1}(E)) = \widehat{f}^{-1}(\gamma_C^p(E)),$$

for every $E \in \Sigma_{M \times Y}$, $C \in \mathcal{C}_M$, and $p \in [0, 1]$.

Finally, the next lemma is a version of [Heifetz & Samet \(1998, Lemma 4.5, p.334\)](#) which takes into account the presence of conditioning events.

Lemma B.2. *Given a product conditional measurable space $(M \times X, \Sigma_M \otimes \Sigma_X, \mathcal{C}_M)$ and an algebra $\mathcal{A}_{M \times X}$ such that $\Sigma_{M \times X} := \sigma(\mathcal{A}_{M \times X})$, if*

$$\mathcal{A}_{\Delta^{\mathcal{C}_M}(M \times X)} := \sigma(\{ \xi_C^p(E) \mid E \in \mathcal{A}_M \otimes \mathcal{A}_X, p \in [0, 1], C \in \mathcal{C}_M \})$$

on $\Delta^{\mathcal{C}_M}(M \times X)$, then $\mathcal{A}_{\Delta^{\mathcal{C}_M}(M \times X)} = \Sigma_{\Delta^{\mathcal{C}_M}(M \times X)}$.

⁷⁹See [Aliprantis & Border \(2006, Lemma 4.10\)](#).


Proof. Observe that the extension of [Definition 2.2](#) to $\Delta^{\mathcal{C}_M}(M \times X)$, i.e., $\Sigma_{\Delta^{\mathcal{C}_M}(M \times X)}$, coincides with the extension of [Definition 2.2](#) to $[\Delta(M \times X)]^{\mathcal{C}_M}$, which is a collection of copies of $\Delta(M \times X)$. Hence, the result follows from [Heifetz & Samet \(1998, Lemma 4.5, p.334\)](#) applied to $[\Delta(M \times X)]^{\mathcal{C}_M} \equiv \Delta(M \times X)$, for every $C \in \mathcal{C}_M$. \blacksquare

C. PROOFS OF SECTION 4

Proof of Lemma 3. We establish the result by proving that

$$\pi_n \circ h^\kappa = \pi_n \circ h^{\kappa'} \circ \mu \tag{C.1}$$

proceeding inductively over $n \in \mathbb{N}$, first by considering the following diagram




and observing that [Equation \(C.1\)](#) is equal to $h_n^\kappa = h_n^{\kappa'} \circ \mu$ from [Definition 4.6](#).

- ($n = 0$) We have that $h_0^{\kappa'} \circ \mu = !_S \circ \mu = !_S = h_0^\kappa$.
- ($n \geq 1$) Assume that the result has been established for n , i.e., $h_n^\kappa = h_n^{\kappa'} \circ \mu$. Thus, we have that

$$\begin{aligned} h_{n+1}^{\kappa'} \circ \mu &= \overline{\mathbf{T}} h_n^{\kappa'} \circ \kappa' \circ \mu \\ &= \overline{\mathbf{T}} h_n^{\kappa'} \circ \overline{\mathbf{T}} \mu \circ \kappa \\ &= \overline{\mathbf{T}} (h_n^{\kappa'} \circ \mu) \circ \kappa \\ &= \overline{\mathbf{T}} h_n^\kappa \circ \kappa \\ &= h_{n+1}^\kappa, \end{aligned}$$

that is made perspicuous by the following diagram



that commutes.

Thus, what is written above establishes the result. \blacksquare

Proof of Lemma 4. We let $\zeta_+ : Z \rightarrow P_{\overline{\mathbf{T}}}$ be defined as $\pi_n^{\overline{\mathbf{T}}} \zeta_+ = \pi_{n+1}$, for every $n \in \mathbb{N}$. We prove that $\pi_n^{\overline{\mathbf{T}}} \zeta_+ \circ \kappa = \pi_n^{\overline{\mathbf{T}}} h_{\overline{\mathbf{T}}}^\kappa \circ \kappa$. Hence, we have that

$$\begin{aligned} \pi_n^{\overline{\mathbf{T}}} \circ h_{\overline{\mathbf{T}}}^\kappa \circ \kappa &= \overline{\mathbf{T}} h_n^\kappa \circ \kappa \\ &= h_{n+1}^\kappa \\ &= \pi_{n+1} \circ h^\kappa \\ &= \pi_n^{\overline{\mathbf{T}}} \circ \zeta_+ \circ h^\kappa. \end{aligned}$$

Thus, this establishes that $\zeta_+ h^\kappa = h_{\mathbf{T}}^\kappa \kappa$ along with the fact that $\zeta_+(Z) \subseteq Z_{\mathbf{T}}$. Concerning the measurability of ζ_+ , we focus on sets belonging to $\Sigma_{\mathbf{T}^{n+1}(\mathbf{1})}$, with $n \in \mathbb{N}$ arbitrary, since the σ -algebra on $Z_{\mathbf{T}}$ is generated by sets of the form $E := (\pi_n^{\mathbf{T}})^{-1}(E_n)$. In light of this point, it is enough to prove that $(\zeta_+)^{-1}(\tilde{E}) \in \Sigma_Z$ for an arbitrary $\tilde{E} := (\pi_n^{\mathbf{T}})^{-1}(\tilde{E}_n)$, which is immediate, since

$$\begin{aligned} (\zeta_+)^{-1}(\tilde{E}) &= (\zeta_+)^{-1}\left((\pi_n^{\mathbf{T}})^{-1}(\tilde{E}_n)\right) \\ &= (\pi_n^{\mathbf{T}} \circ \zeta_+)^{-1}(\tilde{E}_n) \\ &= (\pi_{n+1}^{\mathbf{T}})^{-1}(\tilde{E}_n), \end{aligned}$$

where the set in the last equation is measurable, since $\tilde{E}_n \in \Sigma_{\mathbf{T}^{n+1}(\mathbf{1})}$. \blacksquare

To establish the following lemma, we need three new pieces of notation. First of all, given that

$$\pi_n^{\mathbf{T}} : Z_{\mathbf{T}} \rightarrow \mathbf{T}^{n+1}(\mathbf{1}) := \Delta^{\mathcal{C}_M}(M \times \mathbf{T}^n(\mathbf{1})),$$

we let $\pi_{n,C}^{\mathbf{T}}$ denote the corresponding projection of the function on $[\Delta(M \times \mathbf{T}^n(\mathbf{1}))]^{\{C\}}$, for every $C \in \mathcal{C}_M$. In second place, we let $\nu_C := \nu(\cdot|C)$. Finally, for every $\ell < n$ with $\ell, n \in \mathbb{N}$, let $\rho_{n\ell} : \mathbf{P}_M \mathbf{T}^n(\mathbf{1}) \rightarrow \mathbf{P}_M \mathbf{T}^\ell(\mathbf{1})$ be defined as

$$\rho_{n\ell} := \mathbf{P}_M \mathbf{T}^\ell! \circ \mathbf{P}_M \mathbf{T}^{\ell+1}! \circ \dots \circ \mathbf{P}_M \mathbf{T}^{n-1}!,$$

i.e., the following diagram

$$\begin{array}{ccccccc} \mathbf{P}_M \mathbf{T}^\ell(\mathbf{1}) & \xleftarrow{\mathbf{P}_M \mathbf{T}^\ell!} & \mathbf{P}_M \mathbf{T}^{\ell+1}(\mathbf{1}) & \xleftarrow{\mathbf{P}_M \mathbf{T}^{\ell+1}!} & \dots & \xleftarrow{\mathbf{P}_M \mathbf{T}^{n-1}!} & \mathbf{P}_M \mathbf{T}^n(\mathbf{1}) \\ & & & & \searrow & \swarrow & \\ & & & & \rho_{n\ell} & & \end{array}$$

commutes, with $\rho_{mm} = \mathbf{id}_{\mathbf{P}_M \mathbf{T}^m(\mathbf{1})}$, for every $m \in \mathbb{N}$.

Lemma C.1. *There exists a measurable morphism*

$$\zeta_{\Delta^{\mathcal{C}_M} \mathbf{P}_M} := (\zeta_{\Delta^{\mathcal{C}_M} \mathbf{P}_M})_{C \in \mathcal{C}_M} : Z_{\Delta^{\mathcal{C}_M} \mathbf{P}_M} \rightarrow \Delta^{\mathcal{C}_M}(Z_{\mathbf{P}_M})$$

such that:

- 1) $\zeta_{\Delta^{\mathcal{C}_M} \mathbf{P}_M} \circ h_{\Delta^{\mathcal{C}_M} \mathbf{P}_M}^\kappa = \Delta^{\mathcal{C}_M} h_{\mathbf{P}_M}^\kappa$, for every \mathbf{T} -coalgebra $\mathfrak{C} := \langle S, \kappa \rangle$;
- 2) $\Delta^{\mathcal{C}_M} \pi_n^{\mathbf{P}_M} \circ \zeta_{\Delta^{\mathcal{C}_M}} = \pi_m^{\Delta^{\mathcal{C}_M} \mathbf{P}_M}$, for every $m \in \mathbb{N}$.

Proof. The proof proceeds as follows: first, we define the morphism to then establish its measurability. It should be recalled that $\mathbf{T} := \Delta^{\mathcal{C}_M} \mathbf{P}_M$, a point which we occasionally exploit to lighten the notation.

- *Definition:* Given that $\Sigma_{Z_{\mathbf{P}_M}}$ is generated by the family of sets

$$\mathcal{G} := \left\{ E \mid \exists n \in \mathbb{N}, E_n \in \Sigma_{M \times \mathbf{T}^n(\mathbf{1})} : E = (\pi_n^{\mathbf{P}_M})^{-1}(E_n) \right\},$$

where $E_n \in \Sigma_{M \times \mathbf{T}^n(\mathbf{1})}$ is such that $E = (\pi_n^{\mathbf{P}_M})^{-1}(E_n)$, we define $\zeta_{\Delta^{\mathcal{C}_M} \mathbf{P}_M}$ for the elements of \mathcal{G} to then extend the definition to the elements of $\Sigma_{Z_{\mathbf{P}_M}}$. Thus, let $z \in Z_{\mathbf{T}}$ be arbitrary and let

$$\zeta_{\Delta^{\mathcal{C}_M} \mathbf{P}_M}(z)(E) := \pi_n^{\mathbf{T}}(z)(E_n),$$

i.e., $\zeta_{\Delta^c \mathbf{P}_M}(z)(E) := \pi_{n,C}^{\overline{\mathbf{T}}}(z)(E_n)$, for every $C \in \mathcal{C}_M$ and for every $E_n \in \mathcal{G}$. To see that the definition does not depend on the choice of an $n \in \mathbb{N}$, let $\mathfrak{C} := \langle \mathcal{S}, \kappa \rangle$ be an arbitrary $\overline{\mathbf{T}}$ -coalgebra, let $\nu \in \Delta^c(M \times \mathcal{S})$ be such that $h_{\overline{\mathbf{T}}}^\kappa(\nu) = z$, and let $C \in \mathcal{C}_M$ be arbitrary. Thus, we have that

$$\begin{aligned}
\zeta_{\Delta^c \mathbf{P}_M}(z)(E) &:= \pi_{n,C}^{\overline{\mathbf{T}}}(z)(E_n) \\
&= \pi_n^{\overline{\mathbf{T}}}(h_{\overline{\mathbf{T}}}^\kappa(\nu_C))(E_n) \\
&= (\overline{\mathbf{T}}h_n^\kappa)(\nu_C)(E_n) \\
&= (\Delta^c \mathbf{P}_M h_n^\kappa)(\nu_C)(E_n) \\
&= \nu_C \left((\mathbf{P}_M h_n^\kappa)^{-1}(E_n) \right) \\
&= \nu_C \left((h_{\mathbf{P}_M}^\kappa)^{-1}((\pi_n^{\mathbf{P}_M})^{-1}(E_n)) \right) \\
&= \nu_C \left((h_{\mathbf{P}_M}^\kappa)^{-1}(E) \right) \\
&= (\Delta^c h_{\mathbf{P}_M}^\kappa)(\nu_C)(E),
\end{aligned}$$

where this derivation shows at once that the definition is independent of the natural number $n \in \mathbb{N}$ we started with and that

$$\zeta_{\Delta^c \mathbf{P}_M} h_{\Delta^c \mathbf{P}_M}^\kappa(z)(E) = (\Delta^c h_{\mathbf{P}_M}^\kappa)(\nu_C)(E),$$

for every $E \in \mathcal{G}$. Now, we extend the definition to every $F \in \Sigma_{\mathbf{Z}_{\mathbf{P}_M}}$ by setting

$$\zeta_{\Delta^c \mathbf{P}_M}(z)(F) := (\Delta^c h_{\mathbf{P}_M}^\kappa)(\nu_C)(F), \quad (\text{C.2})$$

for every $C \in \mathcal{C}_M$. We now show that Equation (C.2) is well-defined. Hence, letting $\nu', \nu'' \in \Delta^c(M \times \mathcal{S})$ be arbitrary with $z = h_{\overline{\mathbf{T}}}^\kappa(\nu') = h_{\overline{\mathbf{T}}}^\kappa(\nu'')$, it follows that $\zeta_{\Delta^c \mathbf{P}_M}(h_{\overline{\mathbf{T}}}^\kappa(\nu'))$ and $\zeta_{\Delta^c \mathbf{P}_M}(h_{\overline{\mathbf{T}}}^\kappa(\nu''))$ agree on all the elements of \mathcal{G} . Thus, for every $\ell \leq n$, we have

$$(\pi_n^{\mathbf{P}_M})^{-1}(E_n) \cap (\pi_\ell^{\mathbf{P}_M})^{-1}(E_\ell) = (\pi_n^{\mathbf{P}_M})^{-1}(\rho_{n\ell}^{-1}(E_\ell) \cap E_n).$$

This implies that \mathcal{G} is a π -system, which in turns implies by Lemma B.1 that the CPSs have to agree on all measurable subsets.

- *Measurability:* Recalling Equation (4.10) as a visual aid, for every $E_n \in \Sigma_{M \times \overline{\mathbf{T}}^n(\mathbf{1})}$ and for every $C \in \mathcal{C}_M$, we have that

$$(\Delta^c \mathbf{P}_M \pi_n^{\mathbf{P}_M} \circ \zeta_{\Delta^c \mathbf{P}_M})(z)(E_n) = \zeta_{\Delta^c \mathbf{P}_M}(z) \left((\pi_n^{\mathbf{P}_M})^{-1}(E_n) \right) = \pi_n^{\overline{\mathbf{T}}}(z)(E_n).$$

Hence, from Lemma B.2, it is enough to establish that

$$(\zeta_{\Delta^c \mathbf{P}_M})^{-1} \left((\gamma_C^p(\pi_n^{\mathbf{P}_M})^{-1}(E_n)) \right),$$

for every $C \in \mathcal{C}_M$ and $E_n \in \Sigma_{M \times \overline{\mathbf{T}}^n(\mathbf{1})}$. Thus, we have that

$$\begin{aligned}
(\zeta_{\Delta^c \mathbf{P}_M})^{-1} \left(\gamma_C^p((\pi_n^{\mathbf{P}_M})^{-1}(E_n)) \right) &:= (\zeta_{\Delta^c \mathbf{P}_M})^{-1} \left((\Delta^c \mathbf{P}_M \pi_n^{\mathbf{P}_M})^{-1}(\gamma_C^p(E_n)) \right) \\
&= (\Delta^c \mathbf{P}_M \pi_n^{\mathbf{P}_M} \circ \zeta_{\Delta^c \mathbf{P}_M})^{-1} \left(\gamma_C^p(E_n) \right) \\
&= (\pi_n^{\overline{\mathbf{T}}})^{-1} \left(\gamma_C^p(E_n) \right),
\end{aligned}$$

where the first line is motivated by Remark B.1 and the last set is measurable (with the equality motivated by the fact that we established Condition (2)).

Thus, what is written above establishes the result. ■

Lemma C.2. *There exists a measurable morphism*

$$\zeta_{\mathbf{P}_M} : Z_{\mathbf{K}_M \times \mathbf{Id}} \rightarrow Z_{\mathbf{K}_M} \times Z_{\mathbf{Id}}$$

such that:

- 1) $\zeta_{\mathbf{P}_M} \circ h_{\mathbf{K}_M \times \mathbf{Id}}^\kappa = h_{\mathbf{K}_M}^\kappa \times h_{\mathbf{Id}}^\kappa$, for every $\overline{\mathbf{T}}$ -coalgebra $\mathfrak{C} := \langle S, \kappa \rangle$;
- 2) $(\pi_m^{\mathbf{K}_M} \times \pi_m^{\mathbf{Id}}) \circ \zeta_{\mathbf{P}_M} = \pi_m^{\mathbf{K}_M \times \mathbf{Id}}$, for every $m \in \mathbb{N}$.

Proof. This result is an immediate consequence of [Viglizzo \(2005b, Lemma 6, p.401\)](#) with the binary product applied to the functors \mathbf{K}_M and \mathbf{Id} . \blacksquare

We can now provide the proof of [Lemma 5](#), which turns out to be a straightforward application of the previous lemmata.

Proof of Lemma 5. In light of the fact that $\overline{\mathbf{T}} := \Delta^{\mathfrak{C}_M} \mathbf{P}_M = \Delta^{\mathfrak{C}_M} (\mathbf{K}_M \times \mathbf{Id})$, for every functor $\mathbf{G} \in \{\overline{\mathbf{T}}, \mathbf{P}_M, \mathbf{K}_M, \mathbf{Id}\}$ we define an appropriate measurable morphism $\varphi_{\mathbf{G}}$ satisfying Conditions (1)–(2), i.e., there exists a measurable morphism $\varphi_{\mathbf{G}} : Z_{\mathbf{G}} \rightarrow \mathbf{G}(Z)$ such that:

- 1) $\varphi_{\mathbf{G}} \circ h_{\mathbf{G}}^\kappa = \mathbf{G}h^\kappa$, for every $\overline{\mathbf{T}}$ -coalgebra $\mathfrak{C} := \langle S, \kappa \rangle$;
- 2) $\mathbf{G}\pi_m \circ \varphi_{\mathbf{G}} = \pi_m^{\mathbf{G}}$, for every $m \in \mathbb{N}$.

The morphisms $\varphi_{\mathbf{G}}$ with $\mathbf{G} \in \{\overline{\mathbf{T}}, \mathbf{P}_M, \mathbf{K}_M, \mathbf{Id}\}$ are established in what follows.

- *Functor \mathbf{Id} :* We let $\varphi_{\mathbf{Id}} := \mathbf{id}_Z$, which is measurable and trivially satisfies Conditions (1)–(2), with $\zeta_{\mathbf{Id}} := \varphi_{\mathbf{Id}}$.
- *Functor \mathbf{K}_M :* We let $\varphi_{\mathbf{K}_M} := \pi_0^{\mathbf{K}_M}$, which is measurable.
 - Regarding Condition (1), we let $\mathfrak{C} := \langle S, \kappa \rangle$ be an arbitrary $\overline{\mathbf{T}}$ -coalgebra and we observe that

$$\begin{aligned} \varphi_{\mathbf{K}_M} \circ h_{\mathbf{K}_M}^\kappa &= \pi_0^{\mathbf{K}_M} \circ h_{\mathbf{K}_M}^\kappa \\ &= (\pi_n^{\mathbf{K}_M} \circ h_{\mathbf{K}_M}^\kappa)_{n \in \mathbb{N}} \\ &= \mathbf{K}_M h^\kappa, \end{aligned}$$

thus, establishing the condition.

- Regarding Condition (2), we have that

$$\begin{aligned} \mathbf{K}_M \pi_n \circ \varphi_{\mathbf{K}_M} &= \mathbf{K}_M \pi_n \circ \pi_0^{\mathbf{K}_M} \\ &= \mathbf{id}_M \circ \pi_0^{\mathbf{K}_M} \\ &= \pi_n^{\mathbf{K}_M}. \end{aligned}$$

for every $n \in \mathbb{N}$, thus, establishing the condition.

Finally, we set $\zeta_{\mathbf{K}_M} := \varphi_{\mathbf{K}_M}$.

- *Functor \mathbf{P}_M :* From [Lemma C.2](#), we set

$$\begin{aligned} \varphi_{\mathbf{P}_M} &:= (\pi_0^{\mathbf{K}_M}, \mathbf{id}_Z) \zeta_{\mathbf{P}_M}, \\ &= (\zeta_{\mathbf{K}_M}, \zeta_{\mathbf{Id}}) \zeta_{\mathbf{P}_M}, \end{aligned}$$

which is a measurable morphism that satisfies Conditions (1)–(2) so that the following diagram

$$\begin{array}{ccccc} & & \mathbf{P}_M S = M \times S & & \\ & \swarrow h_{\mathbf{P}_M}^\kappa & \downarrow h_{\mathbf{P}_M}^\kappa & \searrow \mathbf{P}_M h_{\mathbf{Id}}^\kappa & \\ Z_{\mathbf{P}_M} & \xleftarrow{\zeta_{\mathbf{P}_M}} & Z_{\mathbf{K}_M} \times Z_{\mathbf{Id}} & \xrightarrow{(\pi_0^{\mathbf{K}_M}, \mathbf{id}_Z)} & M \times Z \\ & \searrow \pi_n^{\mathbf{P}_M} & \downarrow \pi_n^{\mathbf{P}_M} & \swarrow \mathbf{P}_M \pi_n & \\ & & \mathbf{P}_M \overline{\mathbf{T}}^n(\mathbf{1}) = M \times \overline{\mathbf{T}}^n(\mathbf{1}) & & \end{array}$$

commutes.

- *Functor $\overline{\mathbf{T}}$* : Recalling that $\overline{\mathbf{T}} := \Delta^{\mathbf{c}_M} \mathbf{P}_M$, from [Lemma C.1](#), we set

$$\varphi_{\Delta^{\mathbf{c}_M} \mathbf{P}_M} := (\Delta^{\mathbf{c}_M} \varphi_{\mathbf{P}_M}) \zeta_{\Delta^{\mathbf{c}_M} \mathbf{P}_M},$$

which is a measurable morphism that satisfies Conditions (1)–(2) so that the following diagram

$$\begin{array}{ccccc}
& & \Delta^{\mathbf{c}_M} \mathbf{P}_M \mathbf{S} & & \\
& \swarrow h_{\Delta^{\mathbf{c}_M} \mathbf{P}_M}^\kappa & \downarrow \Delta^{\mathbf{c}_M} h_{\mathbf{P}_M}^\kappa & \searrow \Delta^{\mathbf{c}_M} \mathbf{P}_M h_{\mathbf{Id}}^\kappa & \\
Z_{\Delta^{\mathbf{c}_M} \mathbf{P}_M} & \xrightarrow{\zeta_{\Delta^{\mathbf{c}_M} \mathbf{P}_M}} & \Delta^{\mathbf{c}_M} (Z_{\mathbf{P}_M}) & \xrightarrow{\Delta^{\mathbf{c}_M} \varphi_{\mathbf{P}_M}} & \Delta^{\mathbf{c}_M} (M \times Z) \\
& \searrow \pi_n^{\Delta^{\mathbf{c}_M} \mathbf{P}_M} & \downarrow \Delta^{\mathbf{c}_M} \pi_n^{\mathbf{P}_M} & \swarrow \Delta^{\mathbf{c}_M} \mathbf{P}_M \pi_n & \\
& & \Delta^{\mathbf{c}_M} \mathbf{P}_M \overline{\mathbf{T}}^n(\mathbf{1}) & &
\end{array}$$

commutes as well.

As a result, for every functor $\mathbf{G} \in \{\overline{\mathbf{T}}, \mathbf{P}_M, \mathbf{K}_M, \mathbf{Id}\}$ both Conditions (1)–(2) are satisfied and, in particular, we have that the following diagram

$$\begin{array}{ccccc}
Z_{\Delta^{\mathbf{c}_M} \mathbf{P}_M} & \xrightarrow{\zeta_{\Delta^{\mathbf{c}_M} \mathbf{P}_M}} & \Delta^{\mathbf{c}_M} (Z_{\mathbf{P}_M}) & \xrightarrow{\Delta^{\mathbf{c}_M} \varphi_{\mathbf{P}_M}} & \Delta^{\mathbf{c}_M} (M \times Z) \\
& \searrow & & \searrow & \\
& & & & \varphi_{\Delta^{\mathbf{c}_M} \mathbf{P}_M}
\end{array}$$

commutes, with

$$\begin{aligned}
\vec{\zeta} &:= \varphi_{\Delta^{\mathbf{c}_M} \mathbf{P}_M} \\
&= (\Delta^{\mathbf{c}_M} \varphi_{\mathbf{P}_M}) \zeta_{\Delta^{\mathbf{c}_M} \mathbf{P}_M} \\
&= \left(\Delta^{\mathbf{c}_M} \left((\pi_0^{\mathbf{K}_M}, \mathbf{id}_Z) \zeta_{\mathbf{P}_M} \right) \right) \zeta_{\Delta^{\mathbf{c}_M} \mathbf{P}_M} \\
&= \left(\Delta^{\mathbf{c}_M} \left((\zeta_{\mathbf{K}_M}, \zeta_{\mathbf{Id}}) \zeta_{\mathbf{P}_M} \right) \right) \zeta_{\Delta^{\mathbf{c}_M} \mathbf{P}_M}
\end{aligned}$$

as in [Equation \(4.9\)](#), thus, establishing the result. \blacksquare

Proof of [Lemma 7](#). To establish the result, it is enough to prove that $h_n = \pi_n \circ h = \pi_n$, for every $n \in \mathbb{N}$. Thus, we proceed inductively over $n \in \mathbb{N}$.

- ($n = 0$) We have $h_0 = \pi_0 \circ h = \pi_0 \circ \mathbf{id}_Z$.
- ($n \geq 1$) Assume that the result has been established for n , i.e., $h_n = \pi_n \circ h = \pi_n$. Thus, we have

$$\begin{aligned}
\pi_{n+1} \circ h &= h_{n+1} \\
&= \overline{\mathbf{T}} h_n \circ \zeta \\
&= \overline{\mathbf{T}} \pi_n \circ \zeta \\
&= \overline{\mathbf{T}} \pi_n \circ \vec{\zeta} \circ \zeta_+ \\
&= \pi_n^{\overline{\mathbf{T}}} \circ \zeta_+ \\
&= \pi_{n+1},
\end{aligned}$$

where the second to last equality comes from [Lemma 5](#).

Hence, what is written above establishes the result. \blacksquare

Proof of [Lemma 8](#). Let $\mathfrak{C} := \langle \mathbf{S}, \kappa \rangle$ be an arbitrary $\overline{\mathbf{T}}$ -coalgebra.

- *Existence:* By [Lemma 6](#), h^κ is a $\overline{\mathbf{T}}$ -coalgebra morphism.
- *Uniqueness:* Let $\mu : S \rightarrow Z$ be an arbitrary $\overline{\mathbf{T}}$ -coalgebra morphism. By [Lemma 3](#), we have $h \circ \mu = h^\kappa$ and, from [Lemma 7](#), $h = \mathbf{id}_Z$. Thus, we have that $\mathbf{id}_Z \circ \mu = h^\kappa$, from which it follows that $\mu = h^\kappa$.

Thus, what is written above establishes the result. ■

Proof of [Lemma 10](#). Fix a conditional measurable space $(\Theta, \Sigma_\Theta, \mathcal{C}_\Theta)$ and let $\mathcal{T} := \langle (T_i, \beta_i)_{i \in I} \rangle$ and $\mathcal{T}' := \langle (T'_i, \beta'_i)_{i \in I} \rangle$ be two arbitrary type structures appended to it. From [Lemma 9](#), we can treat them as two \mathbf{T} -coalgebras.

- Regarding Condition (1) in [Definition 3.2](#), the functor \mathbf{P}_Θ ensures that, for every $T \in \mathbf{Ob}(\mathbf{Meas}^I)$, $\mathbf{T}(T)$ implies that there exists a morphism \mathbf{id}_Θ by definition of \mathbf{T} , with $\vartheta_0 = \mathbf{id}_\Theta$.
- Regarding Condition (2) in [Definition 3.2](#), observe that, from [Definition 2.6](#), for a \mathbf{T} -coalgebra morphism $\mu : T \rightarrow T'$, the following diagram

$$\begin{array}{ccc}
 T_i & \xrightarrow{\mu_i} & T'_i \\
 \beta_i \downarrow & & \downarrow \beta'_i \\
 \Delta^{\mathcal{C}_\Theta}(\Theta \times T_{-i}) & \xrightarrow{\widehat{\mu}_{\pm i}} & \Delta^{\mathcal{C}_\Theta}(\Theta \times T'_{-i})
 \end{array}$$

commutes, for every $i \in I$. Clearly, this is exactly Condition (2) in [Definition 3.2](#) with $(\vartheta_i)_{i \in I} = \mu$.

Thus, what is written above establishes the result. ■

REFERENCES

- ACZEL, P. (1988). *Non-Well-Founded Sets*. CSLI. [5]
- ADÁMEK, J.P., KOUBEK, V. (1979). Least Fixed Point of a Functor. *Journal of Computer and System Sciences*, **19**, 163–178. [15]
- (1995). On the Greatest Fixed Point of a Set Functor. *Theoretical Computer Science*, **150**, 57–75. [15]
- ALIPRANTIS, C.D., BORDER, K.C. (2006). *Infinite Dimensional Analysis*. 3rd edition. Springer-Verlag, Berlin. [7, 28, 33]
- ALUFFI, P. (2009). *Algebra: Chapter 0*. American Mathematical Society, Providence. [16]
- ARMBRUSTER, W., BÖGE, W. (1979). Bayesian Game Theory. In *Game theory and related topics*. North-Holland, Amsterdam, (pp. 17–28). [3, 12]
- AUMANN, R.J. (1999b). Interactive Epistemology II: Probability. *International Journal of Game Theory*, **28**, 301–314. [28]
- AUMANN, R.J., HEIFETZ, A. (2002). “Incomplete Information”. In *Handbook of Game Theory with Economic Applications* (R.J. Aumann, S. Hart, editors), volume III. Elsevier/North-Holland, Amsterdam. [3, 28]
- BARR, M. (1993). Terminal Coalgebras in Well-Founded Set Theory. *Theoretical Computer Science*, **114**, 299–315. [15]
- BATTIGALLI, P. (1997). On Rationalizability in Extensive Form Games. *Journal of Economic Theory*, **74**, 40–61. [25]
- BATTIGALLI, P., DI TILLIO, A., GRILLO, E., PENTA, A. (2011). Interactive Epistemology and Solution Concepts for Games with Asymmetric Information. *The B.E. Journal of Theoretical Economics (Advances)*, **11**, Article 6. [5, 23]

- BATTIGALLI, P., FRIEDENBERG, A., SINISCALCHI, M. (Work in Progress). Epistemic Game Theory: Reasoning about Strategic Uncertainty. [3, 24, 29]
- BATTIGALLI, P., SINISCALCHI, M. (1999). Hierarchies of Conditional Beliefs and Interactive Epistemology in Dynamic Games. *Journal of Economic Theory*, **88**, 188–230. [4, 5, 7, 12, 23, 24, 30]
- (2002). Strong Belief and Forward Induction Reasoning. *Journal of Economic Theory*, **106**, 356–391. [24, 25]
- (2003). Rationalization and Incomplete Information. *The B.E. Journal of Theoretical Economics (Advances)*, **3**, Article 3. [25]
- BATTIGALLI, P., TEBALDI, P. (2019). Interactive Epistemology in Simple Dynamic Games with a Continuum of Strategies. *Economic Theory*, **68**, 737–763. [24]
- BEN-PORATH, E. (1997). Rationality, Nash Equilibrium and Backward Induction in Perfect Information Games. *Review of Economic Studies*, **64**, 23–46. [25]
- BERGEMANN, D., MORRIS, S. (2005). Robust Mechanism Design. *Econometrica*, **73**, 1771–1813. [5, 23, 25]
- (2009). Robust Implementation in Direct Mechanisms. *Review of Economic Studies*, **76**, 1175–1204. [25]
- (2011). Robust Implementation in General Mechanisms. *Games and Economic Behavior*, **71**, 261–281. [25]
- (2012). Robust Mechanism Design: An Introduction. In *Robust Mechanism Design. The Role of Private Information and Higher Order Beliefs*. World Scientific. [3]
- BLACKBURN, P., DE RIJKE, M., VENEMA, Y. (2001). *Modal Logic*. Cambridge University Press. [28]
- BOGACHEV, V.I. (2007). *Measure Theory*. Springer-Verlag, Berlin. [7]
- BÖGE, W., EISELE, T. (1979). On Solutions of Bayesian games. *International Journal of Game Theory*, **8**, 193–215. [12]
- BORCEUX, F. (1994). *Handbook of Categorical Algebra 1. Basic Category Theory*. Cambridge University Press. [8]
- BRANDENBURGER, A. (2003). On the Existence of a ‘Complete’ Possibility Structure. In *Cognitive Processes and Economic Behavior* (M. Basili, N. Dimitri, I. Gilboa, editors). Routledge, (pp. 30–34). [12, 27]
- BRANDENBURGER, A., DEKEL, E. (1993). Hierarchies of Beliefs and Common Knowledge. *Journal of Economic Theory*, **59**, 189–198. [3, 4, 29]
- BRANDENBURGER, A., KEISLER, J.H. (2006). An Impossibility Theorem on Beliefs in Games. *Studia Logica: An International Journal for Symbolic Logic*, **84**, 211–240. [12]
- DE OLIVEIRA, H. (2018). Blackwell’s Informativeness Theorem Using Diagrams. *Games and Economic Behavior*, **109**, 126–131. [5]
- DE VITO, N.M. (2023). Complete Conditional Type Structures. *Technical report*, Mimeo. [24]
- DEKEL, E., FUDENBERG, D., MORRIS, S. (2007). Interim Correlated Rationalizability. *Theoretical Economics*, **2**, 15–40. [5, 23]
- DEKEL, E., SINISCALCHI, M. (2015). Epistemic Game Theory. In *Handbook of Game Theory* (H.P. Young, S. Zamir, editors), volume IV. North-Holland, Amsterdam. [3]
- DI TILLIO, A. (2008). Subjective Expected Utility in Games. *Theoretical Economics*, **3**, 287–323. [12]

- DI TILLIO, A., HALPERN, J.Y., SAMET, D. (2014). Conditional Type Spaces. *Games and Economic Behavior*, **87**, 253–268. [29]
- DIÉUDONNÉ, J.A.E. (1948). Sur le théorème de Lebesgue-Nikodym (III). *Annales de l'université de Grenoble*, **23**, 25–53. [28]
- (1989). *A History of Algebraic and Differential Topology, 1900 - 1960*. Birkhäuser, Boston/Basel/Berlin. [1, 29]
- FREYD, P.J. (2008). Algebraic Real Analysis. *Theory and Applications of Categories*, **20**, 215–306. [15]
- FRIEDENBERG, A. (2010). When do Type Structures Contain All Hierarchies of Beliefs? *Games and Economic Behavior*, **68**, 108–129. [12]
- (2019). Bargaining Under Strategic Uncertainty: The Role of Second-Order Optimism. *Econometrica*, **87**, 1835–1865. [25]
- FUKUDA, S. (2021). The Existence of Universal Qualitative Belief Spaces. *Technical report*, Bocconi University & IGIER. [29, 30]
- GALEAZZI, P., MARTI, J. (2023). Choice Structures in Games. *Games and Economic Behavior*, **140**, 431–455. [5]
- GIRY, M. (1982). A Categorical Approach to Probability Theory. In *Categorical Aspects of Topology and Analysis* (B. Banaschewski, editor). Springer Berlin Heidelberg, (pp. 68–85). [4]
- GUARINO, P. (2017). A Topology-Free Construction of the Universal Type Structure for Conditional Probability Systems. *Electronic Proceedings in Theoretical Computer Science*. [1, 30]
- GUO, H., YANNELIS, N.C. (2022). Robust Coalitional Implementation. *Games and Economic Behavior*, **132**, 553–575. [25]
- GUO, Y., SHMAYA, E. (2021). Costly Miscalibration. *Theoretical Economics*, **16**, 477–506. [25]
- HALMOS, P.R. (1950). *Measure Theory*. 1st edition. Litton Educational Publishing, Inc. [28]
- HAMMOND, P.J. (1994). Elementary Non-Archimedean Representations of Probability for Decision Theory and Games. In *Patrick Suppes: Scientific Philosopher* (P. Humphreys, editor), volume 1. Probability and Probabilistic Causality. Springer, Dordrecht. [6]
- HARSANYI, J.C. (1967). Games with Incomplete Information Played by “Bayesian” Players, I–III: Part I. The Basic Model. *Management Science*, **14**, 159–182. [3]
- HEIFETZ, A. (1993). The Bayesian Formulation of Incomplete Information – The Non-Compact Case. *International Journal of Game Theory*, **21**, 329–338. [3]
- (1996). Non-Well-Founded Type Spaces. *Games and Economic Behavior*, **16**, 202–217. [5]
- (1997). Infinitary S5 Epistemic Logic. *Mathematical Logic Quarterly*, **43**, 333–342. [28]
- HEIFETZ, A., MONGIN, P. (2001). Probability Logic for Type Spaces. *Games and Economic Behavior*, **35**, 31–53. [28, 30]
- HEIFETZ, A., SAMET, D. (1998). Topology-Free Typology of Beliefs. *Journal of Economic Theory*, **82**, 324–341. [1, 2, 3, 4, 5, 13, 15, 16, 19, 20, 22, 26, 28, 29, 30, 33, 34]
- (1999). Coherent Beliefs are not always Types. *Journal of Mathematical Economics*, **32**, 475–488. [3, 28]
- HEINSALU, S. (2014). Universal Type Structures with Unawareness. *Games and Economic Behavior*, **83**, 255–266. [2, 5, 27, 29]
- JACOBS, B. (2001). Many-Sorted Coalgebraic Modal Logic: A Model-Theoretic Study. *Theoretical Informatics and Applications*, **35**, 31–59. [16, 29]

- (2017). *Introduction to Coalgebra. Towards Mathematics of States and Observation*. Cambridge University Press. [8, 9, 15, 16, 29]
- JACOBS, B., RUTTEN, J.J.J.M. (1997). A Tutorial on (Co)Algebras and (Co)Induction. *Bulletin of EATCS*, **62**, 222–259. [8, 32]
- JECH, T. (2006). *Set Theory. The Third Millennium Edition, Revised and Expanded*. 4th edition. Springer-Verlag, Berlin. [27]
- KECHRIS, A.S. (1995). *Classical Descriptive Set Theory*. Springer-Verlag, Berlin. [7, 23, 30]
- KURZ, A. (2001). Specifying Coalgebras with Modal Logic. *Theoretical Computer Science*, **260**, 119–138. [29]
- LAMBEK, J. (1968). A Fixpoint Theorem for Complete Categories. *Mathematische Zeitschrift*, **103**, 151–161. [9]
- LAWVERE, F.W. (1962). The Category of Probabilistic Mappings. *Technical report*, Mimeo. [4]
- LEINSTER, T. (2014). *Basic Category Theory*. Cambridge University Press. [5, 8, 16, 27, 32]
- LIU, Q. (2009). On Redundant Types and Bayesian Formulation of Incomplete Information. *Journal of Economic Theory*, **144**, 2115–2145. [12]
- MAC LANE, S. (1998). *Categories for the Working Mathematician*. Springer-Verlag, New York/Berlin/Heidelberg. [8]
- MASCHLER, M., SOLAN, E., ZAMIR, S. (2013). *Game Theory*. Cambridge University Press, Cambridge. [28]
- MEIER, M. (2012). An Infinitary Probability Logic for Type Spaces. *Israel Journal of Mathematics*, **192**, 1–58. [2, 4, 5, 12, 13, 28, 29, 30]
- MEIER, M., PEREA, A. (2023). Forward Induction in a Backward Inductive Manner. *Technical report*, EpiCenter Working Paper. [23, 25]
- MERTENS, J.F., SORIN, S., ZAMIR, S. (2015). *Repeated Games*. Cambridge University Press, Cambridge. [3]
- MERTENS, J.F., ZAMIR, S. (1985). Formulation of Bayesian Analysis for Games with Incomplete Information. *International Journal of Game Theory*, **14**, 1–29. [3, 12, 24, 28, 29]
- MOSS, L. (2011). Connections of Coalgebra and Semantic Modeling. *Proceedings Thirteenth Conference on Theoretical Aspects of Rationality and Knowledge (TARK)*. [5, 14, 28]
- MOSS, L.S., VIGLIZZO, I.D. (2004). Harsanyi Type Spaces and Final Coalgebras Constructed from Satisfied Theories. *Electronic Notes in Theoretical Computer Science*, **106**, 279–295. [5, 13, 16, 28, 30]
- (2006). Final Coalgebras for Functors on Measurable Spaces. *Information and Computation*, **204**, 610–636. [5, 29]
- MÜLLER, C. (2016). Robust Virtual Implementation Under Common Strong Belief in Rationality. *Journal of Economic Theory*, **162**, 407–450. [5, 23]
- MYERSON, R.B. (1986). Multistage Games with Communication. *Econometrica*, **54**, 323–358. [6]
- OSBORNE, M.J., RUBINSTEIN, A. (1994). *A Course in Game Theory*. MIT Press. [24]
- PEARCE, D.G. (1984). Rationalizable Strategic Behavior and the Problem of Perfection. *Econometrica*, **52**, 1029–1050. [25]
- PENTA, A. (2015). Robust Dynamic Implementation. *Journal of Economic Theory*, **160**, 280–316. [5, 23, 25]

- PEREA, A. (2012). *Epistemic Game Theory: Reasoning and Choice*. Cambridge University Press. [3]
- PINTÉR, M. (2010). The Non-Existence of a Universal Topological Type Space. *Journal of Mathematical Economics*, **46**, 223–229. [12]
- PIVATO, M. (Work in Progress). Categorical Decision Theory. [5]
- RÉNYI, A. (1955). On a New Axiomatic Theory of Probability. *Acta Mathematica Academiae Scientiarum Hungarica*, **6**, 285–335. [6, 30]
- RÖßIGER, M. (2001). From Modal Logic to Terminal Coalgebras. *Theoretical Computer Science*, **260**, 209–228. [29]
- SCHUBERT, C. (2009). Terminal Coalgebras for Measure-Polynomial Functors. In *Theory and Applications of Models of Computation* (J. Chen, S.B. Cooper, editors). Springer, Berlin, Heidelberg, (pp. 325–334). [28]
- SCOTT, D.S. (1976). Data Types as Lattices. *SIAM Journal on Computing*, **5**, 522–587. [15]
- SINISCALCHI, M. (2008). “Epistemic Game Theory: Beliefs and Types”. In *The New Palgrave Dictionary of Economics* (S.N. Durlauf, L.E. Blume, editors), 2nd edition. Palgrave Macmillan, New York. [12, 29]
- SMYTH, M.B., PLOTKIN, G.D. (1982). The Category-Theoretic Solution of Recursive Domain Equations. *SIAM Journal on Computing*, **11**, 761–783. [15]
- SPARRE ANDERSEN, E., JESSEN, B. (1948). On the Introduction of Measures in Infinite Product Sets. *Kongelige Danske Videnskabernes Selskab Matematisk-Fysiske Meddelelser*, **25**, 3–7. [28]
- SRIVASTAVA, S.M. (1998). *A Course on Borel Sets*. Springer-Verlag, New York. [6]
- VASSILAKIS, S. (1991). Functional Fixed Points. *Technical Report 33*, Stanford Institute for Theoretical Economics. [12]
- (1992). Some Economic Applications of Scott Domains. *Mathematical Social Sciences*, **24**, 173–208. [12]
- VIGLIZZO, I.D. (2005a). *Coalgebras on measurable spaces*. Ph.D. thesis, Indiana University. [5, 30]
- (2005b). Final Sequences and Final Coalgebras for Measurable Spaces. In *Algebra and Coalgebra in Computer Science* (J.L. Fiadeiro, N. Harman, M. Roggenbach, J. Rutten, editors). Springer Berlin Heidelberg, Berlin, Heidelberg, (pp. 395–407). [2, 4, 5, 13, 15, 16, 18, 26, 27, 28, 29, 33, 37]
- WILSON, R.B. (1987). Game-Theoretic Analyses of Trading Processes. In *Advances in Economics and Econometrics, 5th World Congress of the Econometric Society* (T. Bewley, editor). Cambridge University Press, Cambridge. [3]
- WORRELL, J. (2005). On the Final Sequence of a Finitary Set Functor. *Theoretical Computer Science*, **338**, 184–199. [15]